



EMPLOYEE ILLNESSES AND THE ASSOCIATED EXCLUSIONS/RESTRICTIONS

SALMONELLOSIS	<p>Symptomatic foodhandlers with a confirmed case of Salmonella should be excluded from work and may return only when;</p> <ul style="list-style-type: none"> • the diarrhea has ceased, and • after two consecutive follow-up stool specimens are negative for Salmonella.
SHIGELLOSIS	<p>Symptomatic foodhandlers with a confirmed case of Shigella should be excluded from work and may return only when;</p> <ul style="list-style-type: none"> • the diarrhea has ceased, and • after two consecutive follow-up stool specimens are negative for Shigella.
E.COLI 0157:H7	<p>Foodhandlers with diarrhea should be excluded from work and may return only when;</p> <ul style="list-style-type: none"> • the diarrhea has ceased, and • after two consecutive follow-up stool specimens are negative for E.Coli.
HEPATITIS A	<p>A foodhandler with confirmed Hepatitis A should be excluded from work and may return;</p> <ul style="list-style-type: none"> • 10 days after initial onset of symptoms • Immune globulin (IG) should be administered to restaurant co-workers of the case.
AMEBIASIS	<p>Symptomatic cases should be excluded from work until the symptoms (abdominal pain, diarrhea, fever, drowsiness, and ulcers) resolve.</p> <ul style="list-style-type: none"> • Asymptomatic may return to work or continue to work

VIBRIO CHOLERA	Symptomatic foodhandlers with a case of Cholera should be excluded from work and may return only when; <ul style="list-style-type: none"> • the symptoms have resolved, and • after two consecutive follow-up stool specimens are negative for Cholera.
CRYPTOSPORIDIOSIS	Symptomatic foodhandlers with a case of Cryptosporidiosis should be excluded from work and may return only when; <ul style="list-style-type: none"> • the symptoms have resolved, and • after three consecutive follow-up stool specimens are negative for Cryptosporidiosis.
CYCLOSPORIASIS	Symptomatic foodhandlers infected with Cyclospora should be excluded from work and may return only after diarrhea has ceased AND appropriate antimicrobial therapy has begun.
GIARDIASIS	Symptomatic foodhandlers infected with Giardiasis should be excluded from work and may return only after diarrhea has ceased, and <ul style="list-style-type: none"> • 72 hours of effective antimicrobial therapy have been completed, or • 3 consecutive stools, properly collected, are negative

This exclusion/restriction guide is meant to serve as reference material for the person-in-charge to use in cases of employee illness in a food service operation or retail food establishment. For more information on these or any other illnesses, and to report restrictions