

At A Glance Incidence Reports

The At A Glance reports are meant to be a brief overview of each City of Cleveland neighborhood and every suburban municipality in Cuyahoga County. They show the most recent incidence and mortality rates for each of the 24 different cancer sites/types that could be analyzed at the neighborhood/municipality level. Further details about the individual sections of these reports can be found on the next page.

City of Cleveland Neighborhoods

Brooklyn Centre	Glenville	Old Brooklyn
Buckeye-Shaker	Goodrich-Kirtland Park	Puritas-Longmead
Central	Hough	Riverside
Clark-Fulton	Industrial Valley	South Broadway
Corlett	Jefferson	South Collinwood
Cudell	Kamms Corners	St. Clair-Superior
Detroit-Shoreway/(Near West Side)	Kinsman	Stockyards
Downtown	Lee-Miles	Tremont
Edgewater	Mt. Pleasant	Union-Miles
Euclid-Green	North Broadway	University
Fairfax	North Collinwood	West Boulevard
Forest Hills	Ohio City/Near West Side	Woodland Hills

First Ring Municipalities

Bratenahl	Lakewood
Brooklyn	Linndale
Brooklyn Hts.	Maple Hts.
Brook Park	Newburgh Hts
Cleveland Hts	Parma
Cuyahoga Hts.	Rocky River
East Cleveland	Shaker Hts.
Euclid	South Euclid
Fairview Park	Warrensville Hts.
Garfield Hts.	

Outer Ring Municipalities

Bay Village	Independence	Richmond Hts.
Beachwood	Lyndhurst	Seven Hills.
Bedford	Mayfield Hts.	Solon
Bedford Hts.	Mayfield Village	Strongsville
Bentleyville	Middleburg Hts.	University Hts.
Berea	Moreland Hills	Valley View
Brecksville	North Olmsted	Walton Hills.
Broadview Hts.	North Randall	Westlake
Chagrin Falls Twp.	North Royalton	Woodmere
Chagrin Falls Village	Oakwood	
Gates Mills	Olmsted Falls	
Glenwillow	Olmsted Twp.	
Highland Hills	Orange	
Highland Hts.	Parma Hts	
Hunting Valley	Pepper Pike	

Guide to “At A Glance Incidence Reports”

See Methods/Limitations section for interpretation of calculations for rates, SIRs, and Confidence intervals

City at a Glance: Brooklyn Centre

Name of the community being presented.

Age-adjusted incidence rate for the city, neighborhood, or municipality being presented.

Age-adjusted incidence rate for Cuyahoga County and the State of Ohio for 2002-2006 for comparison.

SEER National age-adjusted incidence rate for 2002-2006

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR)	SIR 95% Confidence Interval	Peer Comparison Rank
Bladder	1.0	24.2	21.5	21.6	21.0	0.97	0.36 – 2.12	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.22	0.25 – 3.55	4
Breast (Female)	4.4	126.8	122.2	120.4	123.8	0.95	0.60 – 1.44	4
Colon & Rectum	5.2	82.1	51.1	52.1	49.1	1.74	1.13 – 2.54	4
Corpus Uterus (female)	1.8	46.6	20.0	26.4	23.3	2.09	0.95 – 3.96	4
Esophagus				5.5	4.5	1.43	0.17 – 5.00	3
Gastric				3.0	2.8	2.50		4

Name of the cancer site or type being presented.

Average annual number of cases is the number diagnosed yearly. To get the total number of cases diagnosed between 2002 and 2006, multiply this number by 5.

The SIR is the observed number of cancer cases divided by the expected. See Methods section for details.

Indicates how this community ranks amongst its peers:
 1- Indicates the community has a rate better than 75% of their peers
 2- Indicates the community has a rate better than 50% of their peers
 3- Indicates the community has a rate better than 25% of their peers
 4- Indicates the community has a rate worse than 75% of their peers

In some instances, peer rankings should be interpreted with some caution because the ranking is based on a rate of cancer that may be unstable due to a small number of cancer cases occurring in the city/neighborhood during the specified time period. These circumstances are easily identified by an average case number being “< 1” and the city rate having an “”.*

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Broadway, South Collinwood

Lists other communities considered to be “peers” of this community, due to similar demographics.

Guide to “At A Glance Incidence Reports”

Additional Information for Understanding the Peer Rankings

The peer rankings are assigned based on the rates for the city/neighborhood. In some instances, peer rankings should be interpreted with some caution because the ranking is based on a rate of cancer that may be unstable due to a small number of cancer cases occurring in the city/neighborhood during the specified time period.

These circumstances are easily identified by an average case number being “< 1” and the city rate having an “*”.

More information on the calculations for the peer rankings can be found in *Appendix II* of the report.

City at a Glance: Cuyahoga County

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	7541.0	477.1	466.4	462.9	1.03	1.01 – 1.04	
Bladder	354.0	21.5	21.6	21.0	1.01	0.97 – 1.06	
Brain and Other CNS	87.2	5.8	6.8	6.4	0.92	0.82 – 0.99	
Breast (Female)	1056.0	122.2	120.4	123.8	0.99	0.96 – 1.01	
Cervix (Female)	61.6	8.0	8.0	8.2	0.96	0.85 – 1.07	
Colon & Rectum	831.4	51.1	52.1	49.1	1.04	1.00 – 1.07	
Corpus Uterus (Female)	258.4	29.9	26.4	23.3	1.28	1.21 – 1.35	
Esophagus	91.4	5.7	5.5	4.5	1.26	1.14 – 1.38	
Hodgkin's Lymphoma	39.6	2.9	3.0	2.8	1.01	0.88 – 1.16	
Kidney & Renal Pelvis	220.4	14.1	14.1	13.6	1.03	0.97 – 1.10	
Larynx	72.0	4.7	4.4	3.5	1.32	1.18 – 1.46	
Leukemia	187.6	11.8	11.1	12.2	0.97	0.91 – 1.03	
Liver & Intrahepatic Bile Duct	86.6	5.6	4.3	6.6	0.83	0.75 – 0.91	
Lung & Bronchus	1141.6	70.7	75.0	63.1	1.10	1.07 – 1.13	
Melanoma of the Skin	234.4	15.3	17.6	19.6	0.78	0.74 – 0.83	
Multiple Myeloma	97.0	6.0	5.1	5.6	1.06	0.97 – 1.16	
Non-Hodgkin's Lymphoma	304.0	19.1	19.2	19.5	0.98	0.93 – 1.03	
Oral Cavity & Pharynx	158.0	10.4	9.6	10.4	0.99	0.92 – 1.06	
Ovary	106.8	12.3	12.3	13.1	0.92	0.84 – 1.00	
Pancreas	199.4	12.3	10.9	11.7	1.04	0.97 – 1.10	
Prostate (Male)	1045.8	157.5	144.7	159.3	0.98	0.95 – 1.01	
Stomach	131.8	8.1	6.1	7.9	1.02	0.94 – 1.10	
Testis (Male)	31.8	4.9	5.5	5.4	0.92	0.78 – 1.07	
Thyroid	114.2	7.9	8.3	9.6	0.82	0.76 – 0.89	

Peer Comparison Neighborhoods/Municipalities:

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Cleveland

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	2174	483.9	477.1	466.4	462.9	1.03	1.01 – 1.05	
Bladder	76.0	16.7	21.5	21.6	21.0	0.79	0.71 – 0.87	
Brain and Other CNS	21.8	4.7	5.8	6.8	6.4	0.73	0.60 – 0.88	
Breast (Female)	283.8	113.3	122.2	120.4	123.8	0.91	0.86 – 0.96	
Cervix (Female)	27.2	11.4	8.0	8.0	8.2	1.35	1.13 – 1.60	
Colon & Rectum	248.8	55	51.1	52.1	49.1	1.11	1.05 – 1.17	
Corpus Uterus (Female)	65.4	26.1	29.9	26.4	23.3	1.11	0.99 – 1.24	
Esophagus	27.0	6.0	5.7	5.5	4.5	1.32	1.11 – 1.56	
Hodgkin's Lymphoma	15.6	3.3	2.9	3.0	2.8	1.20	0.95 – 1.49	
Kidney & Renal Pelvis	65.4	14.7	14.1	14.1	13.6	1.07	0.95 – 1.19	
Larynx	31.2	7.2	4.7	4.4	3.5	2.00	1.70 – 2.34	
Leukemia	41.8	9.1	11.8	11.1	12.2	0.73	0.64 – 0.84	
Liver & Intrahepatic Bile Duct	35.4	8.1	5.6	4.3	6.6	1.19	1.02 – 1.38	
Lung & Bronchus	420.6	93.6	70.7	75.0	63.1	1.45	1.39 – 1.52	
Melanoma of the Skin	27.8	6.1	15.3	17.6	19.6	0.31	0.26 – 0.37	
Multiple Myeloma	29.6	6.5	6.0	5.1	5.6	1.15	0.97 – 1.35	
Non-Hodgkin's Lymphoma	71.0	15.7	19.1	19.2	19.5	0.80	0.72 – 0.88	
Oral Cavity & Pharynx	58.2	13.3	10.4	9.6	10.4	1.26	1.12 – 1.41	
Ovary	28.2	11.0	12.3	12.3	13.1	0.83	0.70 – 0.98	
Pancreas	54.4	12.1	12.3	10.9	11.7	1.02	0.90 – 1.15	
Prostate (Male)	279.2	150.7	157.5	144.7	159.3	0.94	0.89 – 0.99	
Stomach	42.0	9.3	8.1	6.1	7.9	1.16	1.01 – 1.33	
Testis (Male)	7.0	3.0	4.9	5.5	5.4	0.57	0.40 – 0.80	
Thyroid	25.0	5.6	7.9	8.3	9.6	0.57	0.48 – 0.68	

Peer Comparison Neighborhoods/Municipalities:

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: First Ring Suburbs

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	First Ring Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	2538.0	449.3	477.1	466.4	462.9	0.96	0.95 – 0.98	
Bladder	125.0	21.0	21.5	21.6	21.0	0.99	0.91 – 1.07	
Brain and Other CNS	28.2	5.3	5.8	6.8	6.4	0.83	0.70 – 0.98	
Breast (Female)	361	116.2	122.2	120.4	123.8	0.94	0.90 – 0.98	
Cervix (Female)	17.8	6.6	8.0	8.0	8.2	0.78	0.63 – 0.96	
Colon & Rectum	291.8	49.7	51.1	52.1	49.1	1.01	0.96 – 1.06	
Corpus Uterus (Female)	92.8	29.9	29.9	26.4	23.3	1.28	1.16 – 1.40	
Esophagus	30.4	5.4	5.7	5.5	4.5	1.16	0.99 – 1.36	
Hodgkin's Lymphoma	10.2	2.2	2.9	3.0	2.8	0.75	0.56 – 0.99	
Kidney & Renal Pelvis	72.6	13.1	14.1	14.1	13.6	0.96	0.86 – 1.06	
Larynx	20.4	3.7	4.7	4.4	3.5	1.05	0.85 – 1.27	
Leukemia	70.6	12.4	11.8	11.1	12.2	1.02	0.92 – 1.14	
Liver & Intrahepatic Bile Duct	26.0	4.7	5.6	4.3	6.6	0.70	0.59 – 0.83	
Lung & Bronchus	358.2	61.2	70.7	75.0	63.1	0.96	0.91 – 1.00	
Melanoma of the Skin	83.2	15.4	15.3	17.6	19.6	0.78	0.71 – 0.86	
Multiple Myeloma	32.8	5.7	6.0	5.1	5.6	0.99	0.85 – 1.16	
Non-Hodgkin's Lymphoma	104.8	18.4	19.1	19.2	19.5	0.94	0.86 – 1.03	
Oral Cavity & Pharynx	48.4	9.1	10.4	9.6	10.4	0.85	0.75 – 0.97	
Ovary	36.0	11.7	12.3	12.3	13.1	0.86	0.74 – 1.00	
Pancreas	72.6	12.4	12.3	10.9	11.7	1.04	0.94 – 1.16	
Prostate (Male)	350.8	151.9	157.5	144.7	159.3	0.94	0.89 – 0.98	
Stomach	43.8	7.5	8.1	6.1	7.9	0.94	0.82 – 1.07	
Testis (Male)	11.8	5.2	4.9	5.5	5.4	0.99	0.75 – 1.27	
Thyroid	40.4	8.0	7.9	8.3	9.6	0.84	0.73 – 0.96	

Peer Comparison Neighborhoods/Municipalities:

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Outer Ring Suburbs

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	Outer Ring Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	2626.0	462.6	477.1	466.4	462.9	1.00	0.99 – 1.02	
Bladder	144.0	24.0	21.5	21.6	21.0	1.14	1.06 – 1.23	
Brain and Other CNS	35.0	6.8	5.8	6.8	6.4	1.06	0.91 – 1.23	
Breast (Female)	386.2	127.8	122.2	120.4	123.8	1.03	0.99 – 1.08	
Cervix (Female)	14.6	5.7	8.0	8.0	8.2	0.68	0.54 – 0.86	
Colon & Rectum	270.8	45.7	51.1	52.1	49.1	0.94	0.89 – 0.99	
Corpus Uterus (Female)	92.2	30.3	29.9	26.4	23.3	1.31	1.19 – 1.43	
Esophagus	31.6	5.4	5.7	5.5	4.5	1.21	1.03 – 1.41	
Hodgkin's Lymphoma	13.0	2.9	2.9	3.0	2.8	1.04	0.8 – 1.32	
Kidney & Renal Pelvis	78.0	13.9	14.1	14.1	13.6	1.03	0.93 – 1.13	
Larynx	19.2	3.4	4.7	4.4	3.5	0.98	0.79 – 1.20	
Leukemia	70.2	12.7	11.8	11.1	12.2	1.04	0.93 – 1.15	
Liver & Intrahepatic Bile Duct	22.0	3.7	5.6	4.3	6.6	0.59	0.48 – 0.71	
Lung & Bronchus	336.8	56.9	70.7	75.0	63.1	0.91	0.86 – 0.95	
Melanoma of the Skin	115.8	21.6	15.3	17.6	19.6	1.10	1.01 – 1.19	
Multiple Myeloma	31.0	5.3	6.0	5.1	5.6	0.94	0.80 – 1.10	
Non-Hodgkin's Lymphoma	118.8	20.7	19.1	19.2	19.5	1.08	0.99 – 1.17	
Oral Cavity & Pharynx	48.4	8.9	10.4	9.6	10.4	0.85	0.74 – 0.96	
Ovary	39.6	13.0	12.3	12.3	13.1	0.98	0.85 – 1.13	
Pancreas	67.4	11.3	12.3	10.9	11.7	0.97	0.87 – 1.08	
Prostate (Male)	379.4	153.4	157.5	144.7	159.3	0.96	0.92 – 1.00	
Stomach	43.4	7.2	8.1	6.1	7.9	0.94	0.82 – 1.07	
Testis (Male)	11.8	6.0	4.9	5.5	5.4	1.11	0.84 – 1.43	
Thyroid	45.8	9.4	7.9	8.3	9.6	0.98	0.86 – 1.12	

Peer Comparison Neighborhoods/Municipalities:

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Brooklyn Centre

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	35.0	531.9	477.1	466.4	462.9	1.15	0.99 – 1.34	4
Bladder	1.0	24.2	21.5	21.6	21.0	0.97	0.36 – 2.12	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.22	0.25 – 3.55	4
Breast (Female)	4.4	126.8	122.2	120.4	123.8	0.95	0.60 – 1.44	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.58	0.01 – 3.23	1
Colon & Rectum	5.2	82.1	51.1	52.1	49.1	1.74	1.13 – 2.54	4
Corpus Uterus (Female)	1.8	46.6	29.9	26.4	23.3	2.09	0.95 – 3.96	4
Esophagus	<1	*	5.7	5.5	4.5	1.43	0.17 – 5.16	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.50	0.52 – 7.3	4
Kidney & Renal Pelvis	1.0	15.2	14.1	14.1	13.6	1.11	0.36 – 2.58	3
Larynx	<1	*	4.7	4.4	3.5	3.53	0.96 – 9.05	4
Leukemia	<1	*	11.8	11.1	12.2	0.48	0.06 – 1.73	2
Liver & Intrahepatic Bile Duct	1.0	13.6	5.6	4.3	6.6	2.29	0.74 – 5.34	4
Lung & Bronchus	8.0	126.3	70.7	75.0	63.1	2.10	1.50 – 2.86	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.43	0.09 – 1.25	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.73	0.36 – 5.07	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.31	0.04 – 1.13	1
Oral Cavity & Pharynx	1.8	28.1	10.4	9.6	10.4	2.54	1.16 – 4.82	4
Ovary	<1	*	12.3	12.3	13.1	0.41	0.01 – 2.30	1
Pancreas	<1	*	12.3	10.9	11.7	0.58	0.07 – 2.08	1
Prostate (Male)	2.0	79.4	157.5	144.7	159.3	0.47	0.22 – 0.86	2
Stomach	<1	*	8.1	6.1	7.9	0.41	0.01 – 2.29	1
Testis (Male)	<1	*	4.9	5.5	5.4	0.77	0.02 – 4.30	3
Thyroid	<1	*	7.9	8.3	9.6	0.25	0.01 – 1.41	1

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Buckeye-Shaker

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	70.0	439.0	477.1	466.4	462.9	0.93	0.83 – 1.03	1
Bladder	3.0	17.2	21.5	21.6	21.0	0.80	0.44 – 1.35	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.39	0.05 – 1.40	1
Breast (Female)	10.6	117.2	122.2	120.4	123.8	0.90	0.68 – 1.18	3
Cervix (Female)	<1	*	8.0	8.0	8.2	1.05	0.29 – 2.70	3
Colon & Rectum	6.2	38.7	51.1	52.1	49.1	0.76	0.52 – 1.09	1
Corpus Uterus (Female)	1.8	19.4	29.9	26.4	23.3	0.82	0.38 – 1.56	1
Esophagus	<1	*	5.7	5.5	4.5	0.82	0.17 – 2.40	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.44	0.01 – 2.47	1
Kidney & Renal Pelvis	2.2	14	14.1	14.1	13.6	1.01	0.5 – 1.81	1
Larynx	<1	*	4.7	4.4	3.5	0.72	0.09 – 2.61	1
Leukemia	1.0	6.5	11.8	11.1	12.2	0.50	0.16 – 1.16	3
Liver & Intrahepatic Bile Duct	1.0	7.0	5.6	4.3	6.6	0.94	0.31 – 2.20	3
Lung & Bronchus	12.8	79.6	70.7	75.0	63.1	1.24	0.95 – 1.58	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.06	0.00 – 0.35	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.65	0.13 – 1.90	1
Non-Hodgkin's Lymphoma	1.8	11.7	19.1	19.2	19.5	0.56	0.26 – 1.07	2
Oral Cavity & Pharynx	1.4	8.9	10.4	9.6	10.4	0.85	0.34 – 1.75	1
Ovary	1.0	10.7	12.3	12.3	13.1	0.79	0.26 – 1.83	4
Pancreas	1.0	6.4	12.3	10.9	11.7	0.52	0.17 – 1.21	1
Prostate (Male)	14.0	232.7	157.5	144.7	159.3	1.45	1.13 – 1.83	4
Stomach	1.4	8.4	8.1	6.1	7.9	1.07	0.43 – 2.20	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.50	0.01 – 2.79	2
Thyroid	<1	*	7.9	8.3	9.6	0.39	0.08 – 1.14	2

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Central

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	39.0	525.2	477.1	466.4	462.9	1.13	0.98 – 1.30	4
Bladder	1.0	11.4	21.5	21.6	21.0	0.54	0.15 – 1.39	3
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	4.4	90.0	122.2	120.4	123.8	0.84	0.53 – 1.27	2
Cervix (Female)	<1	*	8.0	8.0	8.2	1.50	0.31 – 4.40	3
Colon & Rectum	3.0	40.0	51.1	52.1	49.1	0.86	0.48 – 1.42	1
Corpus Uterus (Female)	1.2	26.1	29.9	26.4	23.3	1.23	0.45 – 2.68	4
Esophagus	<1	*	5.7	5.5	4.5	2.51	0.69 – 6.44	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.06	0.43 – 6.02	4
Kidney & Renal Pelvis	1.4	19.4	14.1	14.1	13.6	1.40	0.56 – 2.88	4
Larynx	1.0	14.2	4.7	4.4	3.5	4.04	1.31 – 9.42	4
Leukemia	<1	*	11.8	11.1	12.2	0.37	0.05 – 1.35	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.65	0.45 – 4.24	3
Lung & Bronchus	12.2	172.3	70.7	75.0	63.1	2.75	2.11 – 3.54	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.01	0.12 – 3.64	2
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.14	0.00 – 0.76	1
Oral Cavity & Pharynx	1.4	21.1	10.4	9.6	10.4	1.84	0.74 – 3.79	3
Ovary	<1	*	12.3	12.3	13.1	0.70	0.08 – 2.52	3
Pancreas	<1	*	12.3	10.9	11.7	0.98	0.27 – 2.50	2
Prostate (Male)	4.2	154	157.5	144.7	159.3	0.89	0.55 – 1.36	2
Stomach	1.0	13.3	8.1	6.1	7.9	1.78	0.58 – 4.15	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.25	0.01 – 1.37	2

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Clark-Fulton

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	43.0	459.9	477.1	466.4	462.9	1.00	0.87 – 1.14	2
Bladder	2.0	24.9	21.5	21.6	21.0	1.11	0.53 – 2.04	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.85	0.17 – 2.47	4
Breast (Female)	5.2	99.1	122.2	120.4	123.8	0.80	0.52 – 1.17	1
Cervix (Female)	1.2	21.6	8.0	8.0	8.2	2.49	0.91 – 5.42	4
Colon & Rectum	4.6	49.6	51.1	52.1	49.1	1.06	0.67 – 1.59	3
Corpus Uterus (Female)	2.2	43	29.9	26.4	23.3	1.81	0.90 – 3.24	4
Esophagus	<1	*	5.7	5.5	4.5	0.50	0.01 – 2.78	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.32	0.63 – 5.95	4
Kidney & Renal Pelvis	1.4	14.4	14.1	14.1	13.6	1.10	0.44 – 2.26	3
Larynx	1.0	10.9	4.7	4.4	3.5	3.15	1.02 – 7.35	4
Leukemia	<1	*	11.8	11.1	12.2	0.16	0.00 – 0.91	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.30	0.35 – 3.32	3
Lung & Bronchus	7.2	79.6	70.7	75.0	63.1	1.30	0.91 – 1.81	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.20	0.02 – 0.74	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.20	0.25 – 3.52	3
Non-Hodgkin's Lymphoma	2.2	23	19.1	19.2	19.5	1.20	0.60 – 2.15	4
Oral Cavity & Pharynx	1.8	18.4	10.4	9.6	10.4	1.82	0.83 – 3.46	3
Ovary	<1	*	12.3	12.3	13.1	0.58	0.07 – 2.08	2
Pancreas	<1	*	12.3	10.9	11.7	0.79	0.22 – 2.03	3
Prostate (Male)	2.8	69.9	157.5	144.7	159.3	0.47	0.26 – 0.79	1
Stomach	<1	*	8.1	6.1	7.9	0.85	0.18 – 2.50	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.74	0.20 – 1.89	3

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Corlett

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	79.0	500.0	477.1	466.4	462.9	1.04	0.94 – 1.14	3
Bladder	2.0	13.7	21.5	21.6	21.0	0.58	0.28 – 1.07	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.19	0.00 – 1.08	2
Breast (Female)	10.8	113.3	122.2	120.4	123.8	0.92	0.69 – 1.20	3
Cervix (Female)	<1	*	8.0	8.0	8.2	1.11	0.30 – 2.85	3
Colon & Rectum	9.6	58.8	51.1	52.1	49.1	1.21	0.90 – 1.61	2
Corpus Uterus (Female)	2.6	27.4	29.9	26.4	23.3	1.14	0.61 – 1.95	3
Esophagus	<1	*	5.7	5.5	4.5	0.53	0.06 – 1.91	2
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	3.0	17.1	14.1	14.1	13.6	1.33	0.75 – 2.2	4
Larynx	<1	*	4.7	4.4	3.5	1.35	0.37 – 3.45	3
Leukemia	1.8	10.9	11.8	11.1	12.2	0.92	0.42 – 1.75	4
Liver & Intrahepatic Bile Duct	1.6	11.1	5.6	4.3	6.6	1.47	0.64 – 2.91	4
Lung & Bronchus	13.0	83.0	70.7	75.0	63.1	1.22	0.94 – 1.55	2
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	1.2	9.0	6.0	5.1	5.6	1.30	0.48 – 2.82	4
Non-Hodgkin's Lymphoma	1.4	8.6	19.1	19.2	19.5	0.45	0.18 – 0.92	2
Oral Cavity & Pharynx	1.2	7.5	10.4	9.6	10.4	0.71	0.26 – 1.55	2
Ovary	<1	*	12.3	12.3	13.1	0.64	0.17 – 1.64	2
Pancreas	2.8	20.1	12.3	10.9	11.7	1.48	0.81 – 2.48	3
Prostate (Male)	17.4	259.5	157.5	144.7	159.3	1.54	1.23 – 1.90	4
Stomach	<1	*	8.1	6.1	7.9	0.31	0.04 – 1.14	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.64	0.02 – 3.55	3
Thyroid	<1	*	7.9	8.3	9.6	0.41	0.09 – 1.21	2

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Cudell

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	31	433.0	477.1	466.4	462.9	0.91	0.77 – 1.06	1
Bladder	1	22.4	21.5	21.6	21.0	1.02	0.41 – 2.09	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.71	0.09 – 2.55	3
Breast (Female)	3.8	95.2	122.2	120.4	123.8	0.73	0.44 – 1.14	1
Cervix (Female)	<1	*	8.0	8.0	8.2	2.05	0.56 – 5.24	4
Colon & Rectum	3.4	46.7	51.1	52.1	49.1	1.01	0.59 – 1.62	1
Corpus Uterus (Female)	1.6	40.4	29.9	26.4	23.3	1.66	0.72 – 3.27	3
Esophagus	<1	*	5.7	5.5	4.5	0.64	0.02 – 3.56	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.42	0.17 – 5.14	3
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.79	0.21 – 2.02	1
Larynx	<1	*	4.7	4.4	3.5	2.38	0.49 – 6.96	2
Leukemia	<1	*	11.8	11.1	12.2	0.84	0.23 – 2.15	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.62	0.44 – 4.15	4
Lung & Bronchus	7.4	105.5	70.7	75.0	63.1	1.73	1.22 – 2.39	3
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.13	0.00 – 0.71	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.03	0.13 – 3.73	2
Non-Hodgkin's Lymphoma	1	14.6	19.1	19.2	19.5	0.69	0.23 – 1.62	3
Oral Cavity & Pharynx	1	15.2	10.4	9.6	10.4	1.26	0.41 – 2.94	3
Ovary	<1	*	12.3	12.3	13.1	0.37	0.01 – 2.04	1
Pancreas	<1	*	12.3	10.9	11.7	0.77	0.16 – 2.26	2
Prostate (Male)	1.8	67.2	157.5	144.7	159.3	0.38	0.17 – 0.72	1
Stomach	0.0	*	8.1	6.1	7.9	0.0	0.00 – 0.00	1
Testis (Male)	<1	*	4.9	5.5	5.4	2.62	0.71 – 6.72	4
Thyroid	<1	*	7.9	8.3	9.6	0.22	0.01 – 1.23	1

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Detroit-Shoreway

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	68.0	482.6	477.1	466.4	462.9	1.04	0.93 – 1.15	2
Bladder	1.0	8.7	21.5	21.6	21.0	0.41	0.15 – 0.88	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.80	0.22 – 2.05	3
Breast (Female)	8.0	104.6	122.2	120.4	123.8	0.83	0.59 – 1.13	2
Cervix (Female)	2.0	26.2	8.0	8.0	8.2	3.05	1.46 – 5.60	4
Colon & Rectum	6.0	43.9	51.1	52.1	49.1	0.86	0.58 – 1.23	1
Corpus Uterus (Female)	1.4	19.7	29.9	26.4	23.3	0.78	0.31 – 1.60	2
Esophagus	1.0	7.1	5.7	5.5	4.5	1.60	0.52 – 3.72	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.30	0.27 – 3.8	3
Kidney & Renal Pelvis	2.2	16.3	14.1	14.1	13.6	1.15	0.57 – 2.05	2
Larynx	1.0	6.9	4.7	4.4	3.5	2.08	0.68 – 4.86	3
Leukemia	1.0	6.3	11.8	11.1	12.2	0.54	0.18 – 1.26	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.86	0.23 – 2.19	1
Lung & Bronchus	14.8	108.9	70.7	75.0	63.1	1.7	1.33 – 2.13	3
Melanoma of the Skin	1.4	9.6	15.3	17.6	19.6	0.49	0.20 – 1.00	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.01	0.28 – 2.60	3
Non-Hodgkin's Lymphoma	3.4	23.2	19.1	19.2	19.5	1.21	0.70 – 1.93	4
Oral Cavity & Pharynx	1.8	12.0	10.4	9.6	10.4	1.22	0.56 – 2.32	2
Ovary	1.6	20.1	12.3	12.3	13.1	1.52	0.66 – 2.99	4
Pancreas	2.0	14.0	12.3	10.9	11.7	1.21	0.58 – 2.23	3
Prostate (Male)	4.2	73.3	157.5	144.7	159.3	0.46	0.29 – 0.71	1
Stomach	1.2	8.5	8.1	6.1	7.9	1.06	0.39 – 2.31	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.85	0.10 – 3.05	4
Thyroid	<1	*	7.9	8.3	9.6	0.53	0.15 – 1.37	3

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Downtown

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	27.0	628.1	477.1	466.4	462.9	1.36	1.14 – 1.61	4
Bladder	1.0	21.5	21.5	21.6	21.0	1.26	0.41 – 2.95	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.64	0.02 – 3.57	3
Breast (Female)	2.4	133.7	122.2	120.4	123.8	1.09	0.56 – 1.90	3
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	2.0	46.1	51.1	52.1	49.1	1.03	0.49 – 1.89	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.95	0.53 – 4.98	4
Esophagus	<1	*	5.7	5.5	4.5	1.13	0.03 – 6.32	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.93	0.6 – 8.57	4
Kidney & Renal Pelvis	1.6	37.9	14.1	14.1	13.6	2.79	1.2 – 5.49	4
Larynx	<1	*	4.7	4.4	3.5	5.69	1.55 – 14.6	4
Leukemia	<1	*	11.8	11.1	12.2	1.18	0.24 – 3.45	4
Liver & Intrahepatic Bile Duct	1.4	31.7	5.6	4.3	6.6	5.07	2.0 – 10.45	4
Lung & Bronchus	3.8	94.6	70.7	75.0	63.1	1.58	0.95 – 2.47	3
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.60	0.12 – 1.77	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	3.62	0.99 – 9.26	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.94	0.26 – 2.40	3
Oral Cavity & Pharynx	1.0	23.2	10.4	9.6	10.4	2.17	0.71 – 5.07	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	3.2	164.8	157.5	144.7	159.3	1.01	0.58 – 1.64	4
Stomach	<1	*	8.1	6.1	7.9	1.25	0.15 – 4.51	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.12	0.14 – 4.04	4
Thyroid	<1	*	7.9	8.3	9.6	0.97	0.20 – 2.84	4

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Edgewater

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	36.0	448.1	477.1	466.4	462.9	0.92	0.80 – 1.07	1
Bladder	2.0	20.6	21.5	21.6	21.0	0.98	0.45 – 1.86	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.11	0.23 – 3.25	4
Breast (Female)	4.6	105.9	122.2	120.4	123.8	0.86	0.55 – 1.29	2
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	3.8	44.7	51.1	52.1	49.1	0.89	0.53 – 1.39	1
Corpus Uterus (Female)	1.0	25.0	29.9	26.4	23.3	1.00	0.33 – 2.34	2
Esophagus	<1	*	5.7	5.5	4.5	0.53	0.01 – 2.97	2
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.18	0 – 1	2
Larynx	<1	*	4.7	4.4	3.5	0.72	0.02 – 4.01	2
Leukemia	<1	*	11.8	11.1	12.2	0.19	0.00 – 1.06	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.48	0.40 – 3.78	3
Lung & Bronchus	9.6	121.9	70.7	75.0	63.1	1.82	1.34 – 2.41	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.23	0.03 – 0.84	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.83	0.10 – 3.01	3
Non-Hodgkin's Lymphoma	1.6	19.2	19.1	19.2	19.5	0.95	0.41 – 1.87	3
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.94	0.26 – 2.40	3
Ovary	<1	*	12.3	12.3	13.1	0.67	0.08 – 2.41	3
Pancreas	2.0	24.4	12.3	10.9	11.7	1.97	0.94 – 3.62	4
Prostate (Male)	4.0	122.9	157.5	144.7	159.3	0.74	0.45 – 1.15	2
Stomach	<1	*	8.1	6.1	7.9	0.58	0.07 – 2.08	2
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.46	0.06 – 1.65	2

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Euclid-Green

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	28.0	530.4	477.1	466.4	462.9	1.21	1.02 – 1.43	3
Bladder	0.0	*	21.5	21.6	21.0	0.41	0.05 – 1.50	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.10	0.13 – 3.97	4
Breast (Female)	4.2	112.3	122.2	120.4	123.8	1.03	0.64 – 1.57	2
Cervix (Female)	<1	*	8.0	8.0	8.2	2.04	0.42 – 5.95	4
Colon & Rectum	3.4	66.7	51.1	52.1	49.1	1.44	0.84 – 2.30	3
Corpus Uterus (Female)	1.2	40.4	29.9	26.4	23.3	1.62	0.60 – 3.53	3
Esophagus	<1	*	5.7	5.5	4.5	1.83	0.22 – 6.60	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.19	0.03 – 6.64	4
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.85	0.18 – 2.49	2
Larynx	<1	*	4.7	4.4	3.5	3.39	0.7 – 9.91	4
Leukemia	<1	*	11.8	11.1	12.2	0.64	0.08 – 2.30	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	2.31	0.63 – 5.92	4
Lung & Bronchus	6.0	119.8	70.7	75.0	63.1	2.03	1.37 – 2.90	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.74	0.02 – 4.10	2
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.61	0.12 – 1.77	2
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	1.07	0.22 – 3.14	2
Ovary	0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	0.37	0.01 – 2.04	1
Prostate (Male)	3.4	158.9	157.5	144.7	159.3	1.12	0.65 – 1.79	4
Stomach	<1	*	8.1	6.1	7.9	2.09	0.57 – 5.36	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	1.02	0.21 – 2.99	4

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Fairfax

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	47.0	540.7	477.1	466.4	462.9	1.08	0.95 – 1.23	4
Bladder	1.0	10.8	21.5	21.6	21.0	0.46	0.15 – 1.06	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.09	0.23 – 3.19	4
Breast (Female)	4.8	116.8	122.2	120.4	123.8	0.79	0.50 – 1.17	3
Cervix (Female)	<1	*	8.0	8.0	8.2	1.21	0.15 – 4.37	2
Colon & Rectum	5.6	60.3	51.1	52.1	49.1	1.13	0.75 – 1.63	4
Corpus Uterus (Female)	1.4	24.0	29.9	26.4	23.3	1.20	0.48 – 2.48	3
Esophagus	1.2	13.0	5.7	5.5	4.5	2.71	0.99 – 5.90	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.99	0.03 – 5.54	3
Kidney & Renal Pelvis	1.6	18.2	14.1	14.1	13.6	1.28	0.55 – 2.52	3
Larynx	1.0	12.7	4.7	4.4	3.5	3.10	1.01 – 7.22	4
Leukemia	<1	*	11.8	11.1	12.2	0.51	0.11 – 1.49	2
Liver & Intrahepatic Bile Duct	1.2	15.0	5.6	4.3	6.6	1.96	0.72 – 4.27	4
Lung & Bronchus	9.8	109.2	70.7	75.0	63.1	1.54	1.14 – 2.03	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	1.4	13.7	6.0	5.1	5.6	2.49	1.00 – 5.13	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.11	0.00 – 0.60	1
Oral Cavity & Pharynx	2.2	30.7	10.4	9.6	10.4	2.38	1.19 – 4.26	4
Ovary	<1	*	12.3	12.3	13.1	0.29	0.01 – 1.63	1
Pancreas	1.8	17.1	12.3	10.9	11.7	1.50	0.68 – 2.84	4
Prostate (Male)	8.6	215.4	157.5	144.7	159.3	1.34	0.97 – 1.81	4
Stomach	1.4	14.9	8.1	6.1	7.9	1.76	0.71 – 3.63	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.35	0.03 – 7.50	4
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Forest Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	84	532.1	477.1	466.4	462.9	1.10	1.00 – 1.21	4
Bladder	1.0	8.6	21.5	21.6	21.0	0.39	0.16 – 0.81	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.38	0.05 – 1.39	2
Breast (Female)	10.2	111.6	122.2	120.4	123.8	0.87	0.65 – 1.14	3
Cervix (Female)	1.0	12.1	8.0	8.0	8.2	1.42	0.46 – 3.32	3
Colon & Rectum	11.2	67.3	51.1	52.1	49.1	1.38	1.05 – 1.80	4
Corpus Uterus (Female)	1.6	15.4	29.9	26.4	23.3	0.71	0.31 – 1.40	1
Esophagus	1.6	10.7	5.7	5.5	4.5	2.10	0.91 – 4.14	4
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	3.0	17.8	14.1	14.1	13.6	1.34	0.75 – 2.21	3
Larynx	1.0	5.1	4.7	4.4	3.5	1.71	0.55 – 3.99	2
Leukemia	2.8	18.4	11.8	11.1	12.2	1.39	0.76 – 2.33	4
Liver & Intrahepatic Bile Duct	1.6	9.6	5.6	4.3	6.6	1.48	0.64 – 2.92	2
Lung & Bronchus	15	94.5	70.7	75.0	63.1	1.37	1.07 – 1.71	2
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	2.2	15.3	6.0	5.1	5.6	2.33	1.16 – 4.17	4
Non-Hodgkin's Lymphoma	3.0	20.1	19.1	19.2	19.5	0.94	0.53 – 1.56	4
Oral Cavity & Pharynx	2.2	15.5	10.4	9.6	10.4	1.33	0.66 – 2.38	3
Ovary	<1	*	12.3	12.3	13.1	0.32	0.04 – 1.15	1
Pancreas	2.2	14.3	12.3	10.9	11.7	1.13	0.57 – 2.03	3
Prostate (Male)	13.4	211.6	157.5	144.7	159.3	1.24	0.96 – 1.57	4
Stomach	2.8	17.2	8.1	6.1	7.9	2.15	1.18 – 3.61	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.56	0.15 – 1.44	3

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Glenville

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	113.0	499.2	477.1	466.4	462.9	1.04	0.96 – 1.13	2
Bladder	2.0	8.3	21.5	21.6	21.0	0.35	0.16 – 0.67	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	15.8	123.9	122.2	120.4	123.8	0.97	0.77 – 1.21	4
Cervix (Female)	1.2	10.0	8.0	8.0	8.2	1.20	0.44 – 2.61	2
Colon & Rectum	15.2	67.8	51.1	52.1	49.1	1.29	1.02 – 1.62	4
Corpus Uterus (Female)	3.6	28.7	29.9	26.4	23.3	1.18	0.70 – 1.86	4
Esophagus	<1	*	5.7	5.5	4.5	0.75	0.20 – 1.92	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.32	0.01 – 1.80	2
Kidney & Renal Pelvis	3.6	16.4	14.1	14.1	13.6	1.15	0.68 – 1.81	3
Larynx	1.4	6.9	4.7	4.4	3.5	1.75	0.7 – 3.6	2
Leukemia	2.2	9.3	11.8	11.1	12.2	0.74	0.37 – 1.33	3
Liver & Intrahepatic Bile Duct	1.8	7.5	5.6	4.3	6.6	1.18	0.54 – 2.25	2
Lung & Bronchus	19.8	85.0	70.7	75.0	63.1	1.30	1.05 – 1.58	1
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	2.0	8.6	6.0	5.1	5.6	1.49	0.71 – 2.74	3
Non-Hodgkin's Lymphoma	3.8	16.1	19.1	19.2	19.5	0.83	0.50 – 1.29	3
Oral Cavity & Pharynx	2.4	10.9	10.4	9.6	10.4	1.02	0.53 – 1.79	1
Ovary	1.0	7.9	12.3	12.3	13.1	0.57	0.18 – 1.33	3
Pancreas	3.2	14.4	12.3	10.9	11.7	1.14	0.65 – 1.84	3
Prostate (Male)	19.4	207.7	157.5	144.7	159.3	1.28	1.04 – 1.56	3
Stomach	2.2	9.3	8.1	6.1	7.9	1.16	0.58 – 2.08	2
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	2.2	10.7	7.9	8.3	9.6	1.07	0.54 – 1.92	4

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Goodrich-Kirtland Park

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	21.0	436.0	477.1	466.4	462.9	0.92	0.75 – 1.11	1
Bladder	1.0	24.7	21.5	21.6	21.0	1.11	0.41 – 2.42	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.33	0.16 – 4.79	4
Breast (Female)	2.8	123.3	122.2	120.4	123.8	0.90	0.49 – 1.51	4
Cervix (Female)	<1	*	8.0	8.0	8.2	4.50	1.2 – 11.52	4
Colon & Rectum	1.8	33.5	51.1	52.1	49.1	0.73	0.33 – 1.38	1
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.30	0.01 – 1.66	1
Larynx	<1	*	4.7	4.4	3.5	1.14	0.03 – 6.34	1
Leukemia	<1	*	11.8	11.1	12.2	0.68	0.08 – 2.44	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.83	0.38 – 5.34	4
Lung & Bronchus	4.8	99.1	70.7	75.0	63.1	1.46	0.94 – 2.18	3
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.21	0.01 – 1.19	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.40	0.17 – 5.06	3
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.62	0.13 – 1.82	3
Oral Cavity & Pharynx	1.4	29.7	10.4	9.6	10.4	2.78	1.12 – 5.72	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	0.34	0.01 – 1.88	1
Prostate (Male)	2.0	93.0	157.5	144.7	159.3	0.55	0.26 – 1.01	2
Stomach	1.0	20.2	8.1	6.1	7.9	2.51	0.82 – 5.87	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Hough

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	91.0	517.7	477.1	466.4	462.9	1.10	1.01 – 1.21	4
Bladder	3.0	14.9	21.5	21.6	21.0	0.72	0.39 – 1.21	4
Brain and Other CNS	1.2	7.2	5.8	6.8	6.4	1.09	0.40 – 2.37	4
Breast (Female)	10.4	106	122.2	120.4	123.8	0.89	0.67 – 1.17	2
Cervix (Female)	1.4	15.8	8.0	8.0	8.2	2.00	0.81 – 4.13	3
Colon & Rectum	9.2	52.5	51.1	52.1	49.1	1.03	0.76 – 1.38	3
Corpus Uterus (Female)	1.4	15.7	29.9	26.4	23.3	0.63	0.25 – 1.31	1
Esophagus	1.0	5.4	5.7	5.5	4.5	1.22	0.40 – 2.86	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.46	0.01 – 2.57	2
Kidney & Renal Pelvis	3.0	18.8	14.1	14.1	13.6	1.26	0.7 – 2.07	4
Larynx	2.2	13.1	4.7	4.4	3.5	3.56	1.78 – 6.37	4
Leukemia	1.6	9.0	11.8	11.1	12.2	0.73	0.31 – 1.44	3
Liver & Intrahepatic Bile Duct	1.2	6.8	5.6	4.3	6.6	1.04	0.38 – 2.25	1
Lung & Bronchus	19.2	109.9	70.7	75.0	63.1	1.65	1.33 – 2.01	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.24	0.07 – 0.62	3
Multiple Myeloma	1.0	5.1	6.0	5.1	5.6	0.98	0.32 – 2.28	2
Non-Hodgkin's Lymphoma	1.8	9.7	19.1	19.2	19.5	0.52	0.24 – 0.99	2
Oral Cavity & Pharynx	2.0	11.8	10.4	9.6	10.4	1.12	0.54 – 2.07	1
Ovary	1.2	12	12.3	12.3	13.1	0.95	0.35 – 2.06	4
Pancreas	2.2	12.9	12.3	10.9	11.7	1.03	0.51 – 1.84	2
Prostate (Male)	14.4	184.9	157.5	144.7	159.3	1.15	0.90 – 1.44	3
Stomach	3.2	17.1	8.1	6.1	7.9	2.23	1.28 – 3.63	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.18	0.14 – 4.26	4
Thyroid	<1	*	7.9	8.3	9.6	0.13	0.00 – 0.74	1

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Industrial Valley

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	4.0	780.4	477.1	466.4	462.9	1.30	0.78 – 2.03	4
Bladder	0.0	*	21.5	21.6	21.0	2.03	0.1 – 11.29	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	<1	*	122.2	120.4	123.8	1.13	0.31 – 2.90	4
Cervix (Female)	<1	*	8.0	8.0	8.2	7.25	1.50 – 21.2	4
Colon & Rectum	<1	*	51.1	52.1	49.1	1.51	0.18 – 5.46	4
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	<1	*	4.7	4.4	3.5	9.73	0.25 – 54.2	4
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	1.0	122.3	70.7	75.0	63.1	3.23	1.05 – 7.54	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	0.0	*	157.5	144.7	159.3	0.00	0.00 – 0.00	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Jefferson

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	90.0	480.2	477.1	466.4	462.9	1.03	0.93 – 1.13	3
Bladder	4.0	20	21.5	21.6	21.0	0.91	0.54 – 1.43	2
Brain and Other CNS	1.4	7.6	5.8	6.8	6.4	1.13	0.45 – 2.32	4
Breast (Female)	11.2	108.4	122.2	120.4	123.8	0.87	0.65 – 1.13	2
Cervix (Female)	1.6	15.8	8.0	8.0	8.2	1.85	0.80 – 3.64	4
Colon & Rectum	9.6	52.1	51.1	52.1	49.1	1.03	0.76 – 1.37	2
Corpus Uterus (Female)	3.8	38	29.9	26.4	23.3	1.58	0.95 – 2.46	4
Esophagus	1.2	6.3	5.7	5.5	4.5	1.43	0.53 – 3.12	3
Hodgkin's Lymphoma	1.2	5.2	2.9	3.0	2.8	2.15	0.79 – 4.67	4
Kidney & Renal Pelvis	3.4	18	14.1	14.1	13.6	1.34	0.78 – 2.14	4
Larynx	<1	*	4.7	4.4	3.5	0.63	0.08 – 2.27	1
Leukemia	2.0	10.7	11.8	11.1	12.2	0.85	0.41 – 1.57	2
Liver & Intrahepatic Bile Duct	1.0	5.6	5.6	4.3	6.6	0.81	0.26 – 1.89	4
Lung & Bronchus	16.6	89.2	70.7	75.0	63.1	1.41	1.12 – 1.74	3
Melanoma of the Skin	3.4	17.3	15.3	17.6	19.6	0.90	0.52 – 1.44	3
Multiple Myeloma	1.2	6.1	6.0	5.1	5.6	1.13	0.42 – 2.47	4
Non-Hodgkin's Lymphoma	3.2	17	19.1	19.2	19.5	0.86	0.49 – 1.40	2
Oral Cavity & Pharynx	2.8	15.7	10.4	9.6	10.4	1.45	0.79 – 2.43	4
Ovary	1.4	12.3	12.3	12.3	13.1	1.01	0.40 – 2.07	4
Pancreas	1.8	9.9	12.3	10.9	11.7	0.82	0.37 – 1.55	1
Prostate (Male)	6.0	79.7	157.5	144.7	159.3	0.50	0.34 – 0.72	1
Stomach	1.0	5.5	8.1	6.1	7.9	0.66	0.22 – 1.55	1
Testis (Male)	<1	*	4.9	5.5	5.4	0.67	0.08 – 2.44	3
Thyroid	<1	*	7.9	8.3	9.6	0.41	0.11 – 1.06	2

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Kamms Corners

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	109.0	461.8	477.1	466.4	462.9	0.98	0.90 – 1.06	2
Bladder	4.0	15.3	21.5	21.6	21.0	0.74	0.45 – 1.14	1
Brain and Other CNS	2.2	9.3	5.8	6.8	6.4	1.55	0.77 – 2.77	4
Breast (Female)	19.6	155.2	122.2	120.4	123.8	1.22	0.99 – 1.49	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.63	0.13 – 1.84	3
Colon & Rectum	11.6	44.9	51.1	52.1	49.1	0.94	0.71 – 1.21	3
Corpus Uterus (Female)	2.4	19.3	29.9	26.4	23.3	0.79	0.41 – 1.38	2
Esophagus	<1	*	5.7	5.5	4.5	0.73	0.20 – 1.86	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.41	0.38 – 3.61	3
Kidney & Renal Pelvis	1.4	6.6	14.1	14.1	13.6	0.44	0.18 – 0.9	1
Larynx	1.2	4.9	4.7	4.4	3.5	1.48	0.54 – 3.22	4
Leukemia	3.0	14.2	11.8	11.1	12.2	1.02	0.57 – 1.69	4
Liver & Intrahepatic Bile Duct	1.2	4.8	5.6	4.3	6.6	0.77	0.28 – 1.68	4
Lung & Bronchus	18.6	77.7	70.7	75.0	63.1	1.18	0.95 – 1.45	4
Melanoma of the Skin	5.4	23.3	15.3	17.6	19.6	1.19	0.79 – 1.74	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.43	0.09 – 1.25	1
Non-Hodgkin's Lymphoma	3.8	15.4	19.1	19.2	19.5	0.80	0.48 – 1.26	2
Oral Cavity & Pharynx	2.4	9.8	10.4	9.6	10.4	1.01	0.52 – 1.76	3
Ovary	2.6	21.9	12.3	12.3	13.1	1.47	0.78 – 2.52	4
Pancreas	2.4	10.8	12.3	10.9	11.7	0.81	0.42 – 1.41	2
Prostate (Male)	11.0	117.4	157.5	144.7	159.3	0.72	0.54 – 0.94	1
Stomach	1.8	7.6	8.1	6.1	7.9	0.90	0.41 – 1.71	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.37	0.01 – 2.05	2
Thyroid	1.4	7.2	7.9	8.3	9.6	0.68	0.27 – 1.40	3

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Kinsman

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	21.0	560.5	477.1	466.4	462.9	1.20	0.98 – 1.45	4
Bladder	0.0	*	21.5	21.6	21.0	0.54	0.06 – 1.93	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.66	0.02 – 3.70	1
Breast (Female)	2.2	110.0	122.2	120.4	123.8	0.82	0.41 – 1.46	3
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	2.0	56.8	51.1	52.1	49.1	1.14	0.55 – 2.10	3
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.60	0.44 – 4.09	4
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.49	0.04 – 8.28	4
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	<1	*	4.7	4.4	3.5	6.32	1.72 – 16.2	4
Leukemia	<1	*	11.8	11.1	12.2	1.12	0.23 – 3.27	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	2.42	0.50 – 7.08	4
Lung & Bronchus	4.0	112.2	70.7	75.0	63.1	1.75	1.07 – 2.71	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.82	0.17 – 2.40	3
Oral Cavity & Pharynx	1.6	42.7	10.4	9.6	10.4	4.16	1.80 – 8.20	4
Ovary	<1	*	12.3	12.3	13.1	0.68	0.02 – 3.81	3
Pancreas	<1	*	12.3	10.9	11.7	1.45	0.30 – 4.24	4
Prostate (Male)	4.0	265.7	157.5	144.7	159.3	1.73	1.06 – 2.67	4
Stomach	<1	*	8.1	6.1	7.9	2.11	0.44 – 6.18	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.50	0.01 – 2.81	4

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Lee-Miles

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	125.0	517.1	477.1	466.4	462.9	1.10	1.02 – 1.19	4
Bladder	4.0	13.5	21.5	21.6	21.0	0.65	0.39 – 1.03	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.45	0.09 – 1.31	1
Breast (Female)	18.4	139.7	122.2	120.4	123.8	1.12	0.91 – 1.38	4
Cervix (Female)	1.0	8.0	8.0	8.0	8.2	1.17	0.38 – 2.73	2
Colon & Rectum	16.4	68.0	51.1	52.1	49.1	1.33	1.06 – 1.65	4
Corpus Uterus (Female)	3.4	26.9	29.9	26.4	23.3	1.05	0.61 – 1.69	2
Esophagus	1.0	3.2	5.7	5.5	4.5	0.86	0.28 – 2.01	1
Hodgkin's Lymphoma	1.6	8.6	2.9	3.0	2.8	3.49	1.51 – 6.87	4
Kidney & Renal Pelvis	2.8	11.6	14.1	14.1	13.6	0.85	0.47 – 1.43	1
Larynx	1.0	5.2	4.7	4.4	3.5	1.14	0.37 – 2.66	3
Leukemia	2.6	10.8	11.8	11.1	12.2	0.92	0.49 – 1.58	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.38	0.08 – 1.10	1
Lung & Bronchus	22.0	81.5	70.7	75.0	63.1	1.31	1.08 – 1.58	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.05	0.00 – 0.26	1
Multiple Myeloma	2.2	8.6	6.0	5.1	5.6	1.53	0.76 – 2.74	4
Non-Hodgkin's Lymphoma	3.8	19.1	19.1	19.2	19.5	0.82	0.49 – 1.27	3
Oral Cavity & Pharynx	1.6	7.6	10.4	9.6	10.4	0.66	0.29 – 1.31	2
Ovary	1.4	10.7	12.3	12.3	13.1	0.79	0.32 – 1.64	2
Pancreas	3.6	16.1	12.3	10.9	11.7	1.20	0.71 – 1.89	3
Prostate (Male)	22.6	214.6	157.5	144.7	159.3	1.29	1.06 – 1.55	3
Stomach	2.6	9.9	8.1	6.1	7.9	1.31	0.70 – 2.25	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.46	0.13 – 1.18	1

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Mt. Pleasant

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	118.0	500.3	477.1	466.4	462.9	1.06	0.98 – 1.15	4
Bladder	3.0	12.3	21.5	21.6	21.0	0.58	0.32 – 0.95	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.53	0.14 – 1.35	3
Breast (Female)	19.4	139.7	122.2	120.4	123.8	1.12	0.91 – 1.37	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.57	0.12 – 1.65	1
Colon & Rectum	12.8	53.7	51.1	52.1	49.1	1.08	0.83 – 1.37	3
Corpus Uterus (Female)	2.4	16.4	29.9	26.4	23.3	0.73	0.38 – 1.28	1
Esophagus	2.0	8.9	5.7	5.5	4.5	1.83	0.88 – 3.36	4
Hodgkin's Lymphoma	1.2	5.6	2.9	3.0	2.8	1.93	0.71 – 4.21	3
Kidney & Renal Pelvis	4.0	17.1	14.1	14.1	13.6	1.24	0.76 – 1.92	4
Larynx	2.2	9.5	4.7	4.4	3.5	2.66	1.33 – 4.75	3
Leukemia	2.4	9.8	11.8	11.1	12.2	0.81	0.42 – 1.41	4
Liver & Intrahepatic Bile Duct	1.4	6.4	5.6	4.3	6.6	0.89	0.36 – 1.84	2
Lung & Bronchus	19.4	82.9	70.7	75.0	63.1	1.24	1.01 – 1.51	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.13	0.03 – 0.39	2
Multiple Myeloma	1.6	6.4	6.0	5.1	5.6	1.17	0.50 – 2.30	3
Non-Hodgkin's Lymphoma	2.0	8.6	19.1	19.2	19.5	0.43	0.21 – 0.79	1
Oral Cavity & Pharynx	2.2	9.5	10.4	9.6	10.4	0.91	0.46 – 1.63	2
Ovary	1.2	9.2	12.3	12.3	13.1	0.65	0.24 – 1.41	3
Pancreas	4.2	17.7	12.3	10.9	11.7	1.47	0.91 – 2.25	4
Prostate (Male)	21.6	234.7	157.5	144.7	159.3	1.46	1.20 – 1.77	4
Stomach	2.4	10.7	8.1	6.1	7.9	1.25	0.65 – 2.19	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.41	0.01 – 2.30	1
Thyroid	1.6	7.3	7.9	8.3	9.6	0.76	0.33 – 1.50	4

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: North Broadway

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	32.0	448.8	477.1	466.4	462.9	0.95	0.81 – 1.11	1
Bladder	2.0	23.3	21.5	21.6	21.0	1.04	0.45 – 2.05	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.77	0.09 – 2.77	3
Breast (Female)	5.0	133.3	122.2	120.4	123.8	1.05	0.68 – 1.55	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.61	0.02 – 3.40	1
Colon & Rectum	3.0	41.0	51.1	52.1	49.1	0.84	0.47 – 1.38	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.68	0.14 – 1.97	2
Esophagus	<1	*	5.7	5.5	4.5	1.22	0.15 – 4.41	2
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	1.8	25.9	14.1	14.1	13.6	1.80	0.82 – 3.42	4
Larynx	<1	*	4.7	4.4	3.5	2.39	0.49 – 6.97	3
Leukemia	<1	*	11.8	11.1	12.2	0.42	0.05 – 1.50	1
Liver & Intrahepatic Bile Duct	1.2	17.9	5.6	4.3	6.6	2.48	0.91 – 5.39	4
Lung & Bronchus	6.8	91.6	70.7	75.0	63.1	1.47	1.02 – 2.05	2
Melanoma of the Skin	1.0	13.9	15.3	17.6	19.6	0.68	0.22 – 1.60	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.49	0.01 – 2.71	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.28	0.03 – 1.00	1
Oral Cavity & Pharynx	2.4	33.7	10.4	9.6	10.4	3.18	1.64 – 5.56	4
Ovary	<1	*	12.3	12.3	13.1	0.77	0.09 – 2.79	4
Pancreas	<1	*	12.3	10.9	11.7	0.71	0.15 – 2.07	1
Prostate (Male)	1.8	53.5	157.5	144.7	159.3	0.34	0.16 – 0.64	1
Stomach	<1	*	8.1	6.1	7.9	0.69	0.08 – 2.50	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.54	0.07 – 1.95	3

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: North Collinwood

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	102.0	470.6	477.1	466.4	462.9	0.99	0.90 – 1.08	2
Bladder	4.0	16.2	21.5	21.6	21.0	0.75	0.45 – 1.17	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.59	0.16 – 1.50	2
Breast (Female)	13.0	111.4	122.2	120.4	123.8	0.86	0.66 – 1.10	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.65	0.13 – 1.90	3
Colon & Rectum	14.0	62.3	51.1	52.1	49.1	1.20	0.94 – 1.52	3
Corpus Uterus (Female)	2.8	23.1	29.9	26.4	23.3	1.00	0.54 – 1.67	2
Esophagus	<1	*	5.7	5.5	4.5	0.79	0.21 – 2.01	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.72	0.09 – 2.59	3
Kidney & Renal Pelvis	2.8	11.8	14.1	14.1	13.6	0.95	0.52 – 1.59	3
Larynx	2.4	12.5	4.7	4.4	3.5	3.24	1.68 – 5.66	3
Leukemia	1.8	8.9	11.8	11.1	12.2	0.64	0.29 – 1.21	3
Liver & Intrahepatic Bile Duct	1.0	4.9	5.6	4.3	6.6	0.70	0.23 – 1.63	2
Lung & Bronchus	18.2	83.4	70.7	75.0	63.1	1.26	1.01 – 1.55	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.19	0.05 – 0.48	2
Multiple Myeloma	1.2	5.8	6.0	5.1	5.6	0.92	0.34 – 2.00	3
Non-Hodgkin's Lymphoma	3.0	13.9	19.1	19.2	19.5	0.68	0.38 – 1.12	2
Oral Cavity & Pharynx	2.6	13.2	10.4	9.6	10.4	1.17	0.63 – 2.01	3
Ovary	2.2	16.4	12.3	12.3	13.1	1.31	0.66 – 2.35	3
Pancreas	3.0	13.2	12.3	10.9	11.7	1.08	0.60 – 1.77	2
Prostate (Male)	13	154.4	157.5	144.7	159.3	0.95	0.73 – 1.21	3
Stomach	1.4	4.6	8.1	6.1	7.9	0.75	0.30 – 1.54	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.79	0.10 – 2.86	4
Thyroid	1.0	4.5	7.9	8.3	9.6	0.51	0.17 – 1.20	3

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Ohio City

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	42.0	509.8	477.1	466.4	462.9	1.12	0.97 – 1.28	3
Bladder	1.0	8.2	21.5	21.6	21.0	0.36	0.08 – 1.06	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.36	0.01 – 2.00	2
Breast (Female)	6.6	159.5	122.2	120.4	123.8	1.26	0.87 – 1.77	4
Cervix (Female)	1.4	34.4	8.0	8.0	8.2	3.92	1.58 – 8.08	4
Colon & Rectum	4.2	50.8	51.1	52.1	49.1	1.08	0.67 – 1.64	2
Corpus Uterus (Female)	1.2	30.3	29.9	26.4	23.3	1.22	0.45 – 2.66	4
Esophagus	<1	*	5.7	5.5	4.5	0.56	0.01 – 3.10	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.57	0.19 – 5.67	4
Kidney & Renal Pelvis	1.0	12.2	14.1	14.1	13.6	0.90	0.29 – 2.1	1
Larynx	1.0	11.7	4.7	4.4	3.5	3.56	1.16 – 8.3	3
Leukemia	1.0	12.8	11.8	11.1	12.2	0.99	0.32 – 2.31	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.47	0.40 – 3.78	3
Lung & Bronchus	8.2	101.5	70.7	75.0	63.1	1.64	1.18 – 2.23	3
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.36	0.07 – 1.06	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.89	0.11 – 3.22	2
Non-Hodgkin's Lymphoma	1.2	13.2	19.1	19.2	19.5	0.75	0.28 – 1.64	3
Oral Cavity & Pharynx	1.6	18.2	10.4	9.6	10.4	1.86	0.80 – 3.67	3
Ovary	1.0	22.2	12.3	12.3	13.1	1.78	0.58 – 4.15	4
Pancreas	1.4	17.8	12.3	10.9	11.7	1.52	0.61 – 3.14	4
Prostate (Male)	3.6	109.6	157.5	144.7	159.3	0.62	0.37 – 0.98	2
Stomach	<1	*	8.1	6.1	7.9	0.95	0.20 – 2.78	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.69	0.14 – 2.02	4

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Old Brooklyn

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	161.0	452.5	477.1	466.4	462.9	0.95	0.88 – 1.02	1
Bladder	10.0	27.3	21.5	21.6	21.0	1.25	0.92 – 1.64	3
Brain and Other CNS	2.0	5.6	5.8	6.8	6.4	0.87	0.42 – 1.61	2
Breast (Female)	18.0	90	122.2	120.4	123.8	0.73	0.59 – 0.90	1
Cervix (Female)	2.0	10.2	8.0	8.0	8.2	1.30	0.62 – 2.39	3
Colon & Rectum	19.4	53.4	51.1	52.1	49.1	1.05	0.85 – 1.28	3
Corpus Uterus (Female)	6.8	38.1	29.9	26.4	23.3	1.46	1.01 – 2.05	4
Esophagus	2.8	7.1	5.7	5.5	4.5	1.69	0.92 – 2.83	4
Hodgkin's Lymphoma	1.4	4.0	2.9	3.0	2.8	1.42	0.57 – 2.93	3
Kidney & Renal Pelvis	4.2	12.0	14.1	14.1	13.6	0.86	0.53 – 1.31	1
Larynx	2.2	6.8	4.7	4.4	3.5	1.78	0.89 – 3.18	4
Leukemia	3.4	9.0	11.8	11.1	12.2	0.75	0.44 – 1.21	1
Liver & Intrahepatic Bile Duct	1.4	4.1	5.6	4.3	6.6	0.59	0.24 – 1.21	2
Lung & Bronchus	31.4	88.4	70.7	75.0	63.1	1.32	1.13 – 1.55	3
Melanoma of the Skin	4.2	12.2	15.3	17.6	19.6	0.59	0.37 – 0.90	2
Multiple Myeloma	1.2	3.1	6.0	5.1	5.6	0.57	0.21 – 1.24	1
Non-Hodgkin's Lymphoma	6.8	19.7	19.1	19.2	19.5	0.94	0.65 – 1.32	3
Oral Cavity & Pharynx	1.8	5.3	10.4	9.6	10.4	0.49	0.22 – 0.93	1
Ovary	3.2	16.8	12.3	12.3	13.1	1.19	0.68 – 1.93	4
Pancreas	5.4	16.0	12.3	10.9	11.7	1.23	0.81 – 1.78	3
Prostate (Male)	14.4	98.7	157.5	144.7	159.3	0.62	0.49 – 0.78	1
Stomach	3.6	9.8	8.1	6.1	7.9	1.21	0.72 – 1.91	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.40	0.05 – 1.43	3
Thyroid	3.0	8.7	7.9	8.3	9.6	0.88	0.49 – 1.45	4

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Puritas-Longmead

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	80.0	552.4	477.1	466.4	462.9	1.16	1.05 – 1.28	3
Bladder	3.0	23.5	21.5	21.6	21.0	1.09	0.64 – 1.75	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.82	0.22 – 2.11	4
Breast (Female)	11.2	139.1	122.2	120.4	123.8	1.11	0.84 – 1.44	4
Cervix (Female)	1.0	13.2	8.0	8.0	8.2	1.52	0.49 – 3.55	4
Colon & Rectum	11.0	75.7	51.1	52.1	49.1	1.52	1.14 – 1.97	4
Corpus Uterus (Female)	4.0	49.1	29.9	26.4	23.3	2.09	1.28 – 3.23	4
Esophagus	1.6	10.7	5.7	5.5	4.5	2.38	1.03 – 4.68	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.43	0.3 – 4.18	4
Kidney & Renal Pelvis	2.2	15.4	14.1	14.1	13.6	1.08	0.54 – 1.93	4
Larynx	1.2	8.4	4.7	4.4	3.5	2.30	0.84 – 5.00	3
Leukemia	1.4	10.0	11.8	11.1	12.2	0.77	0.31 – 1.59	3
Liver & Intrahepatic Bile Duct	1.6	11.3	5.6	4.3	6.6	1.62	0.70 – 3.19	4
Lung & Bronchus	13.8	93.7	70.7	75.0	63.1	1.45	1.13 – 1.84	3
Melanoma of the Skin	1.4	9.6	15.3	17.6	19.6	0.48	0.19 – 0.98	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.72	0.15 – 2.09	2
Non-Hodgkin's Lymphoma	3.6	26.1	19.1	19.2	19.5	1.24	0.73 – 1.96	4
Oral Cavity & Pharynx	2.0	14.0	10.4	9.6	10.4	1.30	0.62 – 2.38	4
Ovary	1.6	20.2	12.3	12.3	13.1	1.49	0.64 – 2.93	4
Pancreas	1.4	9.9	12.3	10.9	11.7	0.81	0.33 – 1.68	2
Prostate (Male)	7.4	121.5	157.5	144.7	159.3	0.72	0.50 – 0.99	2
Stomach	<1	*	8.1	6.1	7.9	0.68	0.19 – 1.75	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.47	0.01 – 2.64	2
Thyroid	1.4	9.0	7.9	8.3	9.6	0.95	0.38 – 1.96	4

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Riverside

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	17.0	377.8	477.1	466.4	462.9	0.80	0.64 – 0.99	1
Bladder	1.0	17.2	21.5	21.6	21.0	0.84	0.23 – 2.15	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.30	0.16 – 4.68	4
Breast (Female)	2.0	80.1	122.2	120.4	123.8	0.63	0.30 – 1.16	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	2.6	59.3	51.1	52.1	49.1	1.17	0.63 – 2.01	2
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.66	0.08 – 2.40	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.32	0.01 – 1.78	2
Larynx	<1	*	4.7	4.4	3.5	1.25	0.03 – 6.97	2
Leukemia	1.6	35.1	11.8	11.1	12.2	2.79	1.20 – 5.49	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.66	0.02 – 3.69	2
Lung & Bronchus	3.2	72.0	70.7	75.0	63.1	1.10	0.63 – 1.78	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.67	0.14 – 1.95	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.56	0.19 – 5.63	3
Non-Hodgkin's Lymphoma	1.4	28.9	19.1	19.2	19.5	1.57	0.63 – 3.24	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.42	0.01 – 2.37	2
Ovary	<1	*	12.3	12.3	13.1	0.59	0.01 – 3.30	2
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	0.0	*	157.5	144.7	159.3	0.26	0.07 – 0.66	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	<1	*	4.9	5.5	5.4	1.7	0.04 – 9.48	4
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: South Broadway

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	91.0	493.0	477.1	466.4	462.9	1.06	0.96 – 1.16	3
Bladder	7.0	36.0	21.5	21.6	21.0	1.69	1.17 – 2.38	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.31	0.04 – 1.13	1
Breast (Female)	10.8	107.7	122.2	120.4	123.8	0.86	0.65 – 1.12	3
Cervix (Female)	<1	*	8.0	8.0	8.2	0.70	0.14 – 2.04	2
Colon & Rectum	8.8	47.0	51.1	52.1	49.1	0.97	0.70 – 1.30	2
Corpus Uterus (Female)	2.6	25.9	29.9	26.4	23.3	1.11	0.59 – 1.90	3
Esophagus	1.8	10.4	5.7	5.5	4.5	2.17	0.99 – 4.13	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.69	0.08 – 2.5	1
Kidney & Renal Pelvis	2.8	14.3	14.1	14.1	13.6	1.11	0.61 – 1.86	2
Larynx	1.4	7.7	4.7	4.4	3.5	2.21	0.89 – 4.56	3
Leukemia	1.2	6.7	11.8	11.1	12.2	0.50	0.19 – 1.10	3
Liver & Intrahepatic Bile Duct	1.0	5.5	5.6	4.3	6.6	0.82	0.27 – 1.91	1
Lung & Bronchus	17.4	95.5	70.7	75.0	63.1	1.49	1.20 – 1.84	3
Melanoma of the Skin	1.0	5.6	15.3	17.6	19.6	0.27	0.09 – 0.62	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.77	0.21 – 1.96	1
Non-Hodgkin's Lymphoma	4.6	23.1	19.1	19.2	19.5	1.25	0.79 – 1.87	4
Oral Cavity & Pharynx	3.4	18.9	10.4	9.6	10.4	1.77	1.03 – 2.83	4
Ovary	2.0	18.9	12.3	12.3	13.1	1.47	0.71 – 2.71	4
Pancreas	2.0	10.4	12.3	10.9	11.7	0.93	0.44 – 1.71	3
Prostate (Male)	9.4	123.4	157.5	144.7	159.3	0.77	0.56 – 1.02	3
Stomach	2.8	15.8	8.1	6.1	7.9	1.90	1.04 – 3.18	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.03	0.21 – 3.00	4
Thyroid	1.6	8.4	7.9	8.3	9.6	0.83	0.36 – 1.64	4

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: South Collinwood

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	54.0	467.2	477.1	466.4	462.9	1.01	0.89 – 1.14	3
Bladder	1.0	13.5	21.5	21.6	21.0	0.60	0.24 – 1.24	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.49	0.06 – 1.76	2
Breast (Female)	6.8	99.7	122.2	120.4	123.8	0.81	0.56 – 1.13	2
Cervix (Female)	<1	*	8.0	8.0	8.2	1.00	0.21 – 2.92	3
Colon & Rectum	6.4	55.6	51.1	52.1	49.1	1.17	0.80 – 1.65	4
Corpus Uterus (Female)	1.4	21.6	29.9	26.4	23.3	0.90	0.36 – 1.85	2
Esophagus	<1	*	5.7	5.5	4.5	1.58	0.43 – 4.04	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.08	0.13 – 3.88	2
Kidney & Renal Pelvis	2.0	16.4	14.1	14.1	13.6	1.27	0.61 – 2.34	4
Larynx	<1	*	4.7	4.4	3.5	0.51	0.01 – 2.82	1
Leukemia	<1	*	11.8	11.1	12.2	0.27	0.03 – 0.98	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.53	0.06 – 1.91	1
Lung & Bronchus	12.4	109.5	70.7	75.0	63.1	1.75	1.34 – 2.25	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.27	0.34 – 3.24	4
Non-Hodgkin's Lymphoma	1.8	15.4	19.1	19.2	19.5	0.8	0.36 – 1.51	3
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.67	0.18 – 1.71	1
Ovary	<1	*	12.3	12.3	13.1	0.67	0.14 – 1.97	3
Pancreas	1.4	12.4	12.3	10.9	11.7	1.08	0.44 – 2.23	4
Prostate (Male)	8.4	192.5	157.5	144.7	159.3	1.14	0.83 – 1.55	3
Stomach	<1	*	8.1	6.1	7.9	0.45	0.05 – 1.63	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.60	0.02 – 3.37	3
Thyroid	<1	*	7.9	8.3	9.6	0.65	0.18 – 1.67	3

Peer Comparison Neighborhoods/Municipalities: Brooklyn Centre, Buckeye-Shaker, Clark-Fulton, Cudell, Mt.Pleasant, South Broadway, South Collinwood

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: St. Clair-Superior

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	42.0	506.4	477.1	466.4	462.9	1.09	0.95 – 1.25	3
Bladder	1.0	6.7	21.5	21.6	21.0	0.36	0.07 – 1.05	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.29	0.35 – 3.29	4
Breast (Female)	4.0	85.3	122.2	120.4	123.8	0.66	0.41 – 1.02	1
Cervix (Female)	<1	*	8.0	8.0	8.2	0.46	0.01 – 2.56	1
Colon & Rectum	4.0	51.3	51.1	52.1	49.1	1.01	0.62 – 1.56	2
Corpus Uterus (Female)	1.2	23.9	29.9	26.4	23.3	1.07	0.39 – 2.33	3
Esophagus	<1	*	5.7	5.5	4.5	0.55	0.01 – 3.04	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	2.0	24.2	14.1	14.1	13.6	1.74	0.84 – 3.21	4
Larynx	<1	*	4.7	4.4	3.5	0.69	0.02 – 3.86	1
Leukemia	1.2	14.4	11.8	11.1	12.2	1.09	0.40 – 2.37	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.72	0.09 – 2.61	1
Lung & Bronchus	7.4	89.2	70.7	75.0	63.1	1.45	1.02 – 2.00	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.35	0.07 – 1.03	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.88	0.11 – 3.17	1
Non-Hodgkin's Lymphoma	2.0	22.8	19.1	19.2	19.5	1.22	0.58 – 2.24	4
Oral Cavity & Pharynx	1.8	22.2	10.4	9.6	10.4	2.05	0.94 – 3.89	4
Ovary	<1	*	12.3	12.3	13.1	0.63	0.08 – 2.26	2
Pancreas	1.4	17.9	12.3	10.9	11.7	1.51	0.61 – 3.11	4
Prostate (Male)	7.8	216.2	157.5	144.7	159.3	1.43	1.02 – 1.96	4
Stomach	<1	*	8.1	6.1	7.9	0.63	0.08 – 2.26	1
Testis (Male)	<1	*	4.9	5.5	5.4	0.81	0.02 – 4.49	4
Thyroid	<1	*	7.9	8.3	9.6	0.44	0.05 – 1.59	3

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Stockyards

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga County Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	33.0	495.8	477.1	466.4	462.9	1.07	0.92 – 1.25	2
Bladder	1.0	20.7	21.5	21.6	21.0	1.08	0.43 – 2.22	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.25	0.26 – 3.66	3
Breast (Female)	3.2	85.5	122.2	120.4	123.8	0.73	0.42 – 1.18	1
Cervix (Female)	1.2	31.7	8.0	8.0	8.2	3.83	1.40 – 8.33	4
Colon & Rectum	3.6	55.2	51.1	52.1	49.1	1.17	0.69 – 1.85	3
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.97	0.26 – 2.47	2
Esophagus	<1	*	5.7	5.5	4.5	1.39	0.17 – 5.01	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.90	0.02 – 5.00	2
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.89	0.24 – 2.27	2
Larynx	<1	*	4.7	4.4	3.5	3.50	0.95 – 8.97	3
Leukemia	<1	*	11.8	11.1	12.2	0.24	0.01 – 1.31	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.92	0.11 – 3.33	1
Lung & Bronchus	9.2	143.1	70.7	75.0	63.1	2.30	1.68 – 3.06	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.45	0.09 – 1.31	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	2.4	38.5	19.1	19.2	19.5	1.87	0.97 – 3.27	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	1.16	0.32 – 2.97	1
Ovary	<1	*	12.3	12.3	13.1	0.43	0.01 – 2.38	2
Pancreas	1.0	15.2	12.3	10.9	11.7	1.38	0.45 – 3.22	4
Prostate (Male)	2.4	85.5	157.5	144.7	159.3	0.52	0.27 – 0.90	1
Stomach	<1	*	8.1	6.1	7.9	0.40	0.01 – 2.24	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Tremont

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	26.0	402.1	477.1	466.4	462.9	0.86	0.72 – 1.02	1
Bladder	1.0	18.7	21.5	21.6	21.0	0.89	0.33 – 1.93	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.8	51.6	122.2	120.4	123.8	0.42	0.19 – 0.80	1
Cervix (Female)	<1	*	8.0	8.0	8.2	2.66	0.72 – 6.81	4
Colon & Rectum	4.6	72.7	51.1	52.1	49.1	1.44	0.91 – 2.16	4
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.5	0.06 – 1.80	1
Esophagus	<1	*	5.7	5.5	4.5	1.38	0.17 – 4.99	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.90	0.02 – 5.02	3
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.90	0.24 – 2.3	1
Larynx	1.0	16.1	4.7	4.4	3.5	4.49	1.5 – 10.49	4
Leukemia	<1	*	11.8	11.1	12.2	0.94	0.26 – 2.42	4
Liver & Intrahepatic Bile Duct	1.2	18.4	5.6	4.3	6.6	2.78	1.02 – 6.05	4
Lung & Bronchus	5.0	78.3	70.7	75.0	63.1	1.24	0.80 – 1.83	1
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	2.20	0.60 – 5.63	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.61	0.17 – 1.57	2
Oral Cavity & Pharynx	1.0	16.1	10.4	9.6	10.4	1.47	0.48 – 3.42	3
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	0.27	0.01 – 1.48	1
Prostate (Male)	1.4	52.4	157.5	144.7	159.3	0.31	0.13 – 0.65	1
Stomach	<1	*	8.1	6.1	7.9	1.16	0.24 – 3.39	2
Testis (Male)	<1	*	4.9	5.5	5.4	2.53	0.52 – 7.40	4
Thyroid	<1	*	7.9	8.3	9.6	0.28	0.01 – 1.57	2

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Union-Miles

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	70.0	515.1	477.1	466.4	462.9	1.09	0.98 – 1.21	3
Bladder	1.0	7.2	21.5	21.6	21.0	0.29	0.08 – 0.74	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.85	0.23 – 2.18	3
Breast (Female)	7.6	89.7	122.2	120.4	123.8	0.75	0.53 – 1.02	1
Cervix (Female)	<1	*	8.0	8.0	8.2	1.22	0.33 – 3.12	2
Colon & Rectum	8.2	62.1	51.1	52.1	49.1	1.27	0.91 – 1.72	4
Corpus Uterus (Female)	2.0	24.5	29.9	26.4	23.3	1.02	0.49 – 1.88	3
Esophagus	<1	*	5.7	5.5	4.5	0.96	0.20 – 2.82	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.51	0.31 – 4.42	4
Kidney & Renal Pelvis	2.0	14.1	14.1	14.1	13.6	1.05	0.5 – 1.93	2
Larynx	<1	*	4.7	4.4	3.5	0.80	0.10 – 2.89	1
Leukemia	1.2	9.8	11.8	11.1	12.2	0.71	0.26 – 1.55	3
Liver & Intrahepatic Bile Duct	1.6	11.9	5.6	4.3	6.6	1.73	0.75 – 3.41	3
Lung & Bronchus	14.2	103.5	70.7	75.0	63.1	1.63	1.28 – 2.06	3
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.15	0.02 – 0.54	3
Multiple Myeloma	1.2	9.9	6.0	5.1	5.6	1.58	0.58 – 3.44	3
Non-Hodgkin's Lymphoma	2.2	16.4	19.1	19.2	19.5	0.83	0.42 – 1.49	4
Oral Cavity & Pharynx	2.2	15.4	10.4	9.6	10.4	1.51	0.76 – 2.71	2
Ovary	<1	*	12.3	12.3	13.1	0.37	0.05 – 1.35	2
Pancreas	1.6	12.3	12.3	10.9	11.7	1.05	0.45 – 2.06	2
Prostate (Male)	12	199	157.5	144.7	159.3	1.30	0.99 – 1.67	3
Stomach	1.6	16.3	8.1	6.1	7.9	1.54	0.66 – 3.03	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	1.0	7.4	7.9	8.3	9.6	0.74	0.24 – 1.72	4

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: University

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	37.0	448.7	477.1	466.4	462.9	0.79	0.68 – 0.92	1
Bladder	1.0	8.4	21.5	21.6	21.0	0.47	0.17 – 1.03	2
Brain and Other CNS	1.0	23.5	5.8	6.8	6.4	1.70	0.55 – 3.96	4
Breast (Female)	4.0	107.1	122.2	120.4	123.8	0.64	0.39 – 0.99	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.62	0.02 – 3.43	2
Colon & Rectum	4.4	47.2	51.1	52.1	49.1	0.78	0.49 – 1.18	2
Corpus Uterus (Female)	1.6	24.5	29.9	26.4	23.3	1.36	0.59 – 2.68	3
Esophagus	<1	*	5.7	5.5	4.5	1.29	0.27 – 3.77	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.6	0.02 – 3.35	3
Kidney & Renal Pelvis	2.2	32.0	14.1	14.1	13.6	1.77	0.88 – 3.16	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.44	0.09 – 1.28	3
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.98	0.20 – 2.87	2
Lung & Bronchus	6.4	71.5	70.7	75.0	63.1	0.95	0.65 – 1.34	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.11	0.00 – 0.61	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.98	0.20 – 2.87	2
Non-Hodgkin's Lymphoma	1.2	11.4	19.1	19.2	19.5	0.58	0.21 – 1.27	2
Oral Cavity & Pharynx	1.0	12.1	10.4	9.6	10.4	1.08	0.35 – 2.52	2
Ovary	<1	*	12.3	12.3	13.1	1.05	0.28 – 2.68	3
Pancreas	1.2	12.1	12.3	10.9	11.7	0.87	0.32 – 1.89	2
Prostate (Male)	5.6	160.1	157.5	144.7	159.3	1.01	0.67 – 1.46	2
Stomach	1.0	7.2	8.1	6.1	7.9	1.10	0.36 – 2.57	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.61	0.02 – 3.39	3
Thyroid	<1	*	7.9	8.3	9.6	0.78	0.16 – 2.29	4

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: West Boulevard

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	64.0	467.4	477.1	466.4	462.9	1.03	0.92 – 1.14	2
Bladder	4.0	31.3	21.5	21.6	21.0	1.50	0.91 – 2.31	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.82	0.22 – 2.11	3
Breast (Female)	7.6	102.3	122.2	120.4	123.8	0.82	0.58 – 1.12	1
Cervix (Female)	1.0	11.5	8.0	8.0	8.2	1.49	0.48 – 3.48	4
Colon & Rectum	7.2	55.7	51.1	52.1	49.1	1.13	0.79 – 1.57	2
Corpus Uterus (Female)	1.4	19.2	29.9	26.4	23.3	0.81	0.33 – 1.67	1
Esophagus	1.0	7.5	5.7	5.5	4.5	1.71	0.55 – 3.99	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.29	0.27 – 3.78	4
Kidney & Renal Pelvis	1.8	11.4	14.1	14.1	13.6	0.98	0.45 – 1.86	3
Larynx	<1	*	4.7	4.4	3.5	0.87	0.11 – 3.14	2
Leukemia	1.2	8.8	11.8	11.1	12.2	0.70	0.26 – 1.52	2
Liver & Intrahepatic Bile Duct	1.4	11.1	5.6	4.3	6.6	1.58	0.63 – 3.25	3
Lung & Bronchus	13.8	102.6	70.7	75.0	63.1	1.69	1.32 – 2.14	3
Melanoma of the Skin	2.0	13.4	15.3	17.6	19.6	0.71	0.34 – 1.31	4
Multiple Myeloma	1.2	8.4	6.0	5.1	5.6	1.64	0.60 – 3.57	4
Non-Hodgkin's Lymphoma	3.0	21.3	19.1	19.2	19.5	1.13	0.63 – 1.87	4
Oral Cavity & Pharynx	1.8	13.5	10.4	9.6	10.4	1.27	0.58 – 2.42	4
Ovary	<1	*	12.3	12.3	13.1	0.40	0.05 – 1.46	2
Pancreas	<1	*	12.3	10.9	11.7	0.40	0.08 – 1.18	2
Prostate (Male)	5.0	90.8	157.5	144.7	159.3	0.57	0.37 – 0.84	2
Stomach	2.0	15.0	8.1	6.1	7.9	1.94	0.93 – 3.56	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.81	0.10 – 2.92	3
Thyroid	<1	*	7.9	8.3	9.6	0.53	0.14 – 1.35	3

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Woodland Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	45.0	507.3	477.1	466.4	462.9	1.09	0.95 – 1.24	3
Bladder	1.0	10.5	21.5	21.6	21.0	0.57	0.18 – 1.32	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.62	0.07 – 2.23	2
Breast (Female)	8.2	155.5	122.2	120.4	123.8	1.24	0.89 – 1.69	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.87	0.11 – 3.14	1
Colon & Rectum	4.4	51.1	51.1	52.1	49.1	1.06	0.66 – 1.60	2
Corpus Uterus (Female)	1.8	33.6	29.9	26.4	23.3	1.45	0.66 – 2.75	4
Esophagus	1.4	17.0	5.7	5.5	4.5	3.57	1.44 – 7.36	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.07	0.43 – 6.04	4
Kidney & Renal Pelvis	1.4	15.8	14.1	14.1	13.6	1.15	0.46 – 2.37	2
Larynx	<1	*	4.7	4.4	3.5	1.29	0.16 – 4.65	2
Leukemia	<1	*	11.8	11.1	12.2	0.35	0.04 – 1.25	1
Liver & Intrahepatic Bile Duct	1.2	13.5	5.6	4.3	6.6	2.04	0.75 – 4.43	3
Lung & Bronchus	8.2	94.8	70.7	75.0	63.1	1.50	1.08 – 2.04	2
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	1.2	14.2	6.0	5.1	5.6	2.49	0.91 – 5.41	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.23	0.03 – 0.84	1
Oral Cavity & Pharynx	1.2	12.6	10.4	9.6	10.4	1.29	0.47 – 2.81	2
Ovary	<1	*	12.3	12.3	13.1	0.29	0.01 – 1.60	2
Pancreas	<1	*	12.3	10.9	11.7	0.41	0.05 – 1.47	1
Prostate (Male)	6.2	181.8	157.5	144.7	159.3	1.10	0.75 – 1.56	3
Stomach	<1	*	8.1	6.1	7.9	1.19	0.32 – 3.05	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.22	0.01 – 1.20	2

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Bay Village

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	86.0	466.0	477.1	466.4	462.9	0.98	0.89 – 1.08	4
Bladder	4.0	21.1	21.5	21.6	21.0	1.01	0.62 – 1.56	2
Brain and Other CNS	1.0	6.2	5.8	6.8	6.4	0.87	0.28 – 2.04	3
Breast (Female)	12.6	127.8	122.2	120.4	123.8	0.97	0.75 – 1.24	3
Cervix (Female)	<1	*	8.0	8.0	8.2	0.26	0.01 – 1.43	3
Colon & Rectum	8.2	45.0	51.1	52.1	49.1	0.88	0.63 – 1.20	1
Corpus Uterus (Female)	2.4	24.0	29.9	26.4	23.3	0.98	0.51 – 1.72	1
Esophagus	<1	*	5.7	5.5	4.5	0.70	0.14 – 2.04	2
Hodgkin's Lymphoma	1.2	6.4	2.9	3.0	2.8	2.78	1.02 – 6.06	4
Kidney & Renal Pelvis	2.2	11.8	14.1	14.1	13.6	0.85	0.43 – 1.53	2
Larynx	<1	*	4.7	4.4	3.5	0.90	0.19 – 2.63	4
Leukemia	3.0	16.3	11.8	11.1	12.2	1.36	0.76 – 2.24	3
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.16	0.00 – 0.87	1
Lung & Bronchus	8.4	44.7	70.7	75.0	63.1	0.70	0.51 – 0.95	1
Melanoma of the Skin	6.6	38.9	15.3	17.6	19.6	1.83	1.26 – 2.57	4
Multiple Myeloma	1.0	5.0	6.0	5.1	5.6	0.93	0.30 – 2.18	3
Non-Hodgkin's Lymphoma	4.0	20.9	19.1	19.2	19.5	1.10	0.67 – 1.70	2
Oral Cavity & Pharynx	2.4	13.7	10.4	9.6	10.4	1.21	0.63 – 2.12	4
Ovary	1.4	12.2	12.3	12.3	13.1	1.03	0.41 – 2.11	3
Pancreas	1.4	7.6	12.3	10.9	11.7	0.64	0.26 – 1.31	2
Prostate (Male)	13.8	164.3	157.5	144.7	159.3	1.04	0.81 – 1.31	4
Stomach	1.2	6.3	8.1	6.1	7.9	0.81	0.30 – 1.76	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.59	0.33 – 4.66	3
Thyroid	1.0	4.8	7.9	8.3	9.6	0.59	0.19 – 1.38	1

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Beachwood

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	118.0	454.4	477.1	466.4	462.9	0.99	0.91 – 1.07	2
Bladder	7.0	19.5	21.5	21.6	21.0	1.02	0.71 – 1.42	2
Brain and Other CNS	1.8	14.5	5.8	6.8	6.4	1.43	0.65 – 2.72	4
Breast (Female)	15.8	139.4	122.2	120.4	123.8	1.00	0.79 – 1.25	3
Cervix (Female)	1.0	9.2	8.0	8.0	8.2	1.45	0.47 – 3.38	4
Colon & Rectum	14.0	41.3	51.1	52.1	49.1	0.94	0.74 – 1.19	3
Corpus Uterus (Female)	3.8	35.2	29.9	26.4	23.3	1.27	0.77 – 1.99	3
Esophagus	1.0	2.9	5.7	5.5	4.5	0.81	0.26 – 1.89	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.54	0.01 – 3.02	3
Kidney & Renal Pelvis	4.4	17.7	14.1	14.1	13.6	1.35	0.85 – 2.04	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	4.2	16.1	11.8	11.1	12.2	1.29	0.80 – 1.96	3
Liver & Intrahepatic Bile Duct	1.2	2.8	5.6	4.3	6.6	0.74	0.27 – 1.62	3
Lung & Bronchus	16.2	54.0	70.7	75.0	63.1	0.91	0.72 – 1.13	3
Melanoma of the Skin	5.2	23.4	15.3	17.6	19.6	1.21	0.79 – 1.77	1
Multiple Myeloma	1.4	4.5	6.0	5.1	5.6	0.86	0.35 – 1.78	3
Non-Hodgkin's Lymphoma	6.8	26.1	19.1	19.2	19.5	1.32	0.92 – 1.85	3
Oral Cavity & Pharynx	1.4	7.9	10.4	9.6	10.4	0.59	0.24 – 1.22	3
Ovary	1.6	9.4	12.3	12.3	13.1	0.87	0.37 – 1.71	3
Pancreas	2.4	5.1	12.3	10.9	11.7	0.66	0.34 – 1.15	1
Prostate (Male)	13.2	135.6	157.5	144.7	159.3	0.81	0.63 – 1.03	2
Stomach	3.0	9.7	8.1	6.1	7.9	1.26	0.71 – 2.08	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.00	0.03 – 5.55	3
Thyroid	1.6	10.6	7.9	8.3	9.6	1.15	0.50 – 2.27	3

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Bedford

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	89.0	504.7	477.1	466.4	462.9	1.05	0.96 – 1.15	4
Bladder	4.0	22.8	21.5	21.6	21.0	0.95	0.58 – 1.47	2
Brain and Other CNS	1.0	7.0	5.8	6.8	6.4	0.94	0.31 – 2.20	4
Breast (Female)	10.8	115.8	122.2	120.4	123.8	0.90	0.68 – 1.18	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.86	0.18 – 2.52	4
Colon & Rectum	9.8	53.5	51.1	52.1	49.1	1.03	0.76 – 1.36	3
Corpus Uterus (Female)	2.4	25.2	29.9	26.4	23.3	1.06	0.55 – 1.85	2
Esophagus	2.2	13.1	5.7	5.5	4.5	2.60	1.30 – 4.66	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.43	0.29 – 4.17	4
Kidney & Renal Pelvis	3.4	20.3	14.1	14.1	13.6	1.41	0.82 – 2.25	4
Larynx	1.2	6.1	4.7	4.4	3.5	1.95	0.72 – 4.25	4
Leukemia	2.0	11.8	11.8	11.1	12.2	0.89	0.43 – 1.64	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.68	0.19 – 1.74	3
Lung & Bronchus	17.0	94.7	70.7	75.0	63.1	1.40	1.12 – 1.73	4
Melanoma of the Skin	2.0	12.2	15.3	17.6	19.6	0.59	0.28 – 1.08	2
Multiple Myeloma	1.0	6.3	6.0	5.1	5.6	0.93	0.30 – 2.16	3
Non-Hodgkin's Lymphoma	3.6	19.5	19.1	19.2	19.5	1.00	0.59 – 1.58	3
Oral Cavity & Pharynx	1.6	9.9	10.4	9.6	10.4	0.89	0.39 – 1.76	3
Ovary	<1	*	12.3	12.3	13.1	0.45	0.09 – 1.32	1
Pancreas	2.4	11.2	12.3	10.9	11.7	1.04	0.54 – 1.82	3
Prostate (Male)	13	180.2	157.5	144.7	159.3	1.09	0.84 – 1.39	3
Stomach	2.0	10.0	8.1	6.1	7.9	1.30	0.62 – 2.40	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	1.0	6.2	7.9	8.3	9.6	0.67	0.22 – 1.56	2

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Bedford Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	59.0	484.5	477.1	466.4	462.9	1.03	0.91 – 1.15	4
Bladder	2.0	21.2	21.5	21.6	21.0	0.79	0.38 – 1.46	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.52	0.06 – 1.87	1
Breast (Female)	8.4	119.6	122.2	120.4	123.8	0.95	0.69 – 1.29	3
Cervix (Female)	<1	*	8.0	8.0	8.2	0.36	0.01 – 2.00	2
Colon & Rectum	6.6	54.3	51.1	52.1	49.1	1.11	0.77 – 1.56	4
Corpus Uterus (Female)	1.0	13.2	29.9	26.4	23.3	0.60	0.19 – 1.39	1
Esophagus	<1	*	5.7	5.5	4.5	1.07	0.22 – 3.11	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.23	0.15 – 4.45	3
Kidney & Renal Pelvis	2.0	19.1	14.1	14.1	13.6	1.17	0.56 – 2.15	4
Larynx	<1	*	4.7	4.4	3.5	0.90	0.11 – 3.23	3
Leukemia	1.2	8.7	11.8	11.1	12.2	0.84	0.31 – 1.83	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.48	0.06 – 1.72	2
Lung & Bronchus	8.8	73.6	70.7	75.0	63.1	1.12	0.81 – 1.50	4
Melanoma of the Skin	1.2	9.1	15.3	17.6	19.6	0.50	0.18 – 1.08	1
Multiple Myeloma	1.2	9.1	6.0	5.1	5.6	1.73	0.64 – 3.77	4
Non-Hodgkin's Lymphoma	1.6	12.8	19.1	19.2	19.5	0.67	0.29 – 1.33	1
Oral Cavity & Pharynx	1.2	9.4	10.4	9.6	10.4	0.91	0.34 – 1.99	2
Ovary	1.2	16.0	12.3	12.3	13.1	1.31	0.48 – 2.85	4
Pancreas	<1	*	12.3	10.9	11.7	0.43	0.09 – 1.25	1
Prostate (Male)	13.4	245.3	157.5	144.7	159.3	1.52	1.18 – 1.94	4
Stomach	<1	*	8.1	6.1	7.9	0.84	0.23 – 2.15	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.69	0.02 – 3.85	3
Thyroid	<1	*	7.9	8.3	9.6	0.67	0.18 – 1.72	2

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Bentleyville

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	4.0	776.6	477.1	466.4	462.9	1.07	0.63 – 1.69	4
Bladder	0.0	*	21.5	21.6	21.0	0.00	0.00 – 0.00	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	3.70	0.09 – 20.6	4
Breast (Female)	<1	*	122.2	120.4	123.8	1.07	0.22 – 3.14	4
Cervix (Female)	<1	*	8.0	8.0	8.2	9.94	1.2 – 35.91	4
Colon & Rectum	<1	*	51.1	52.1	49.1	1.26	0.15 – 4.54	4
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	3.74	0.45 – 13.5	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.51	0.01 – 2.81	1
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	1.43	0.04 – 7.97	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	2.25	0.06 – 12.6	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	2.81	0.07 – 15.7	4
Prostate (Male)	0.0	*	157.5	144.7	159.3	1.11	0.23 – 3.24	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Berea

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	96.0	473.5	477.1	466.4	462.9	1.02	0.93 – 1.12	3
Bladder	7.0	32.2	21.5	21.6	21.0	1.54	1.07 – 2.15	4
Brain and Other CNS	1.0	5.3	5.8	6.8	6.4	0.80	0.26 – 1.87	2
Breast (Female)	14.4	131.5	122.2	120.4	123.8	1.04	0.81 – 1.31	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.72	0.15 – 2.12	3
Colon & Rectum	8.6	41.1	51.1	52.1	49.1	0.84	0.61 – 1.14	2
Corpus Uterus (Female)	4.6	39.5	29.9	26.4	23.3	1.76	1.11 – 2.64	4
Esophagus	1.6	7.5	5.7	5.5	4.5	1.74	0.75 – 3.42	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.36	0.01 – 2.01	1
Kidney & Renal Pelvis	3.0	15.2	14.1	14.1	13.6	1.11	0.62 – 1.82	2
Larynx	<1	*	4.7	4.4	3.5	0.58	0.07 – 2.08	2
Leukemia	2.2	10.7	11.8	11.1	12.2	0.89	0.44 – 1.59	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.30	0.04 – 1.09	2
Lung & Bronchus	12.6	61.9	70.7	75.0	63.1	0.97	0.74 – 1.24	3
Melanoma of the Skin	5.6	27.7	15.3	17.6	19.6	1.44	0.96 – 2.08	4
Multiple Myeloma	1.4	6.3	6.0	5.1	5.6	1.21	0.48 – 2.48	3
Non-Hodgkin's Lymphoma	5.8	28.3	19.1	19.2	19.5	1.46	0.98 – 2.09	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.19	0.02 – 0.70	1
Ovary	1.4	13.3	12.3	12.3	13.1	0.92	0.37 – 1.90	4
Pancreas	1.8	8.5	12.3	10.9	11.7	0.74	0.34 – 1.40	1
Prostate (Male)	12	149.9	157.5	144.7	159.3	0.92	0.70 – 1.18	2
Stomach	1	4.9	8.1	6.1	7.9	0.61	0.20 – 1.42	1
Testis (Male)	<1	*	4.9	5.5	5.4	0.80	0.10 – 2.88	4
Thyroid	2.6	13.9	7.9	8.3	9.6	1.42	0.76 – 2.43	4

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Bratenahl

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	11.0	528.9	477.1	466.4	462.9	0.99	0.75 – 1.28	3
Bladder	0.0	*	21.5	21.6	21.0	0.35	0.01 – 1.97	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.8	147.7	122.2	120.4	123.8	1.21	0.56 – 2.31	3
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.2	44.2	51.1	52.1	49.1	0.94	0.34 – 2.04	3
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.41	0.17 – 5.09	4
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.60	0.02 – 3.34	2
Larynx	<1	*	4.7	4.4	3.5	4.50	0.6 – 16.26	4
Leukemia	<1	*	11.8	11.1	12.2	0.72	0.02 – 4.00	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.21	0.03 – 6.71	4
Lung & Bronchus	1.0	42.0	70.7	75.0	63.1	0.59	0.19 – 1.38	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	1.82	0.50 – 4.67	3
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	1.68	0.46 – 4.30	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	1.60	0.19 – 5.79	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	1.29	0.16 – 4.65	3
Prostate (Male)	1.6	137.4	157.5	144.7	159.3	0.77	0.33 – 1.52	2
Stomach	<1	*	8.1	6.1	7.9	0.98	0.02 – 5.44	3
Testis (Male)	<1	*	4.9	5.5	5.4	7.02	0.18 – 39.1	4
Thyroid	1.0	65.7	7.9	8.3	9.6	5.70	1.85 – 13.3	4

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Brecksville

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	71.0	397.3	477.1	466.4	462.9	0.85	0.77 – 0.94	1
Bladder	3.0	17.1	21.5	21.6	21.0	0.81	0.47 – 1.32	2
Brain and Other CNS	1.2	6.7	5.8	6.8	6.4	1.15	0.42 – 2.51	3
Breast (Female)	12.4	137.5	122.2	120.4	123.8	1.11	0.85 – 1.42	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.93	0.19 – 2.71	4
Colon & Rectum	5.8	32.2	51.1	52.1	49.1	0.64	0.43 – 0.92	1
Corpus Uterus (Female)	2.8	31.2	29.9	26.4	23.3	1.33	0.73 – 2.23	3
Esophagus	1.0	5.3	5.7	5.5	4.5	1.19	0.39 – 2.79	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.64	0.34 – 4.79	4
Kidney & Renal Pelvis	1.4	7.9	14.1	14.1	13.6	0.57	0.23 – 1.17	1
Larynx	<1	*	4.7	4.4	3.5	0.62	0.08 – 2.25	2
Leukemia	1.8	10.4	11.8	11.1	12.2	0.86	0.39 – 1.63	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.49	0.10 – 1.45	3
Lung & Bronchus	8.2	45.1	70.7	75.0	63.1	0.69	0.50 – 0.94	2
Melanoma of the Skin	3.6	21.5	15.3	17.6	19.6	1.07	0.63 – 1.69	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.76	0.21 – 1.96	3
Non-Hodgkin's Lymphoma	1.6	8.6	19.1	19.2	19.5	0.46	0.2 – 0.90	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.21	0.03 – 0.77	1
Ovary	<1	*	12.3	12.3	13.1	0.51	0.10 – 1.48	2
Pancreas	2.8	15.1	12.3	10.9	11.7	1.29	0.70 – 2.16	4
Prostate (Male)	14.4	156.2	157.5	144.7	159.3	0.98	0.77 – 1.24	3
Stomach	1.0	5.5	8.1	6.1	7.9	0.69	0.22 – 1.60	3
Testis (Male)	<1	*	4.9	5.5	5.4	2.71	0.74 – 6.93	4
Thyroid	1.0	5.9	7.9	8.3	9.6	0.67	0.22 – 1.57	1

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Broadview Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	82.0	458.1	477.1	466.4	462.9	0.97	0.88 – 1.07	3
Bladder	5.0	27.4	21.5	21.6	21.0	1.28	0.83 – 1.89	4
Brain and Other CNS	1.4	7.8	5.8	6.8	6.4	1.26	0.51 – 2.59	3
Breast (Female)	12.4	127.2	122.2	120.4	123.8	1.02	0.79 – 1.31	3
Cervix (Female)	<1	*	8.0	8.0	8.2	1.06	0.29 – 2.72	4
Colon & Rectum	8.8	49.8	51.1	52.1	49.1	0.98	0.71 – 1.31	4
Corpus Uterus (Female)	2.0	21.8	29.9	26.4	23.3	0.88	0.42 – 1.62	1
Esophagus	<1	*	5.7	5.5	4.5	0.97	0.26 – 2.47	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.90	0.11 – 3.26	3
Kidney & Renal Pelvis	2.2	12.4	14.1	14.1	13.6	0.89	0.45 – 1.6	3
Larynx	<1	*	4.7	4.4	3.5	0.63	0.08 – 2.29	2
Leukemia	2	11.7	11.8	11.1	12.2	0.92	0.44 – 1.70	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.50	0.10 – 1.46	4
Lung & Bronchus	8.8	47.2	70.7	75.0	63.1	0.75	0.54 – 1.01	1
Melanoma of the Skin	4.2	24.1	15.3	17.6	19.6	1.21	0.75 – 1.84	3
Multiple Myeloma	1.2	6.8	6.0	5.1	5.6	1.16	0.42 – 2.51	3
Non-Hodgkin's Lymphoma	2.2	12.1	19.1	19.2	19.5	0.62	0.31 – 1.11	1
Oral Cavity & Pharynx	3.2	19.0	10.4	9.6	10.4	1.72	0.98 – 2.80	4
Ovary	1.2	12.4	12.3	12.3	13.1	0.93	0.34 – 2.03	3
Pancreas	1.4	8.1	12.3	10.9	11.7	0.65	0.26 – 1.34	2
Prostate (Male)	13.4	173.3	157.5	144.7	159.3	1.04	0.81 – 1.32	4
Stomach	1.0	5.3	8.1	6.1	7.9	0.69	0.22 – 1.61	2
Testis (Male)	<1	*	4.9	5.5	5.4	1.99	0.54 – 5.10	4
Thyroid	2.2	13.0	7.9	8.3	9.6	1.34	0.67 – 2.39	4

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Brooklyn

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	78.0	468.9	477.1	466.4	462.9	0.99	0.90 – 1.09	3
Bladder	6.0	35.1	21.5	21.6	21.0	1.58	1.07 – 2.24	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.85	0.23 – 2.17	2
Breast (Female)	10.6	125.9	122.2	120.4	123.8	0.97	0.73 – 1.27	3
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	7.6	44.5	51.1	52.1	49.1	0.86	0.61 – 1.19	2
Corpus Uterus (Female)	3.8	43.1	29.9	26.4	23.3	1.81	1.09 – 2.83	4
Esophagus	<1	*	5.7	5.5	4.5	1.00	0.27 – 2.56	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.57	0.01 – 3.18	2
Kidney & Renal Pelvis	2.4	14.9	14.1	14.1	13.6	1.06	0.55 – 1.85	2
Larynx	1.0	6.6	4.7	4.4	3.5	1.70	0.55 – 3.96	4
Leukemia	1.6	10	11.8	11.1	12.2	0.80	0.34 – 1.57	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.72	0.20 – 1.85	4
Lung & Bronchus	10.8	59.3	70.7	75.0	63.1	0.93	0.70 – 1.21	2
Melanoma of the Skin	2.4	16.1	15.3	17.6	19.6	0.78	0.41 – 1.37	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.79	0.22 – 2.02	1
Non-Hodgkin's Lymphoma	4.0	24.1	19.1	19.2	19.5	1.21	0.74 – 1.87	4
Oral Cavity & Pharynx	1.4	9.6	10.4	9.6	10.4	0.84	0.34 – 1.74	3
Ovary	<1	*	12.3	12.3	13.1	0.50	0.10 – 1.47	1
Pancreas	3.2	19.0	12.3	10.9	11.7	1.50	0.86 – 2.44	4
Prostate (Male)	9.4	124.2	157.5	144.7	159.3	0.79	0.58 – 1.05	1
Stomach	1.4	7.7	8.1	6.1	7.9	0.99	0.40 – 2.04	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.36	0.16 – 4.91	4
Thyroid	1.4	9.5	7.9	8.3	9.6	1.10	0.44 – 2.26	4

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Brooklyn Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	12.0	639.6	477.1	466.4	462.9	1.28	0.98 – 1.64	4
Bladder	1.0	32.6	21.5	21.6	21.0	1.31	0.27 – 3.82	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.6	146.5	122.2	120.4	123.8	1.20	0.52 – 2.36	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.2	64.0	51.1	52.1	49.1	1.16	0.42 – 2.51	4
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.57	0.19 – 5.68	4
Esophagus	<1	*	5.7	5.5	4.5	2.08	0.05 – 11.6	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	4.55	0.12 – 25.4	4
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	1.43	0.17 – 5.18	4
Larynx	0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.83	0.02 – 4.61	2
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	2.0	82.2	70.7	75.0	63.1	1.45	0.69 – 2.66	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.53	0.01 – 2.93	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	3.32	0.4 – 11.98	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	1.50	0.31 – 4.38	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.97	0.02 – 5.39	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	0.80	0.02 – 4.46	3
Prostate (Male)	1.8	187.4	157.5	144.7	159.3	1.22	0.56 – 2.32	4
Stomach	1.0	47.1	8.1	6.1	7.9	5.99	1.95 – 14.0	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	3.62	0.75 – 10.6	4

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Brook Park

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	139.0	551.1	477.1	466.4	462.9	1.14	1.06 – 1.23	4
Bladder	8.0	35.1	21.5	21.6	21.0	1.52	1.10 – 2.06	4
Brain and Other CNS	1.8	8.8	5.8	6.8	6.4	1.16	0.53 – 2.20	4
Breast (Female)	14.8	105.7	122.2	120.4	123.8	0.85	0.67 – 1.07	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.78	0.21 – 2.00	2
Colon & Rectum	13.2	52.4	51.1	52.1	49.1	1.04	0.80 – 1.32	2
Corpus Uterus (Female)	5.0	37.6	29.9	26.4	23.3	1.47	0.95 – 2.17	3
Esophagus	1.2	4.4	5.7	5.5	4.5	0.99	0.36 – 2.15	2
Hodgkin's Lymphoma	1.0	4.8	2.9	3.0	2.8	1.65	0.54 – 3.86	3
Kidney & Renal Pelvis	3.2	12.4	14.1	14.1	13.6	0.89	0.51 – 1.45	1
Larynx	1.6	6.1	4.7	4.4	3.5	1.67	0.72 – 3.29	3
Leukemia	3.6	14.3	11.8	11.1	12.2	1.21	0.71 – 1.91	4
Liver & Intrahepatic Bile Duct	1.2	4.2	5.6	4.3	6.6	0.69	0.25 – 1.50	3
Lung & Bronchus	27.2	104.5	70.7	75.0	63.1	1.58	1.32 – 1.87	4
Melanoma of the Skin	7.4	30.4	15.3	17.6	19.6	1.52	1.07 – 2.09	4
Multiple Myeloma	1.6	5.7	6.0	5.1	5.6	1.08	0.46 – 2.12	3
Non-Hodgkin's Lymphoma	5.2	24.3	19.1	19.2	19.5	1.04	0.68 – 1.53	4
Oral Cavity & Pharynx	2.8	11.4	10.4	9.6	10.4	1.04	0.57 – 1.74	3
Ovary	1.6	11.9	12.3	12.3	13.1	0.87	0.38 – 1.71	3
Pancreas	4.0	16.3	12.3	10.9	11.7	1.32	0.80 – 2.03	4
Prostate (Male)	17.8	154.2	157.5	144.7	159.3	0.89	0.72 – 1.10	2
Stomach	2.4	9.2	8.1	6.1	7.9	1.18	0.61 – 2.06	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.13	0.23 – 3.30	4
Thyroid	2.4	10.3	7.9	8.3	9.6	1.06	0.55 – 1.85	4

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Chagrin Falls Township

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	1.0	945.5	477.1	466.4	462.9	1.88	0.69 – 4.10	4
Bladder	0.0	*	21.5	21.6	21.0	0.00	0.00 – 0.00	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	0.0	*	122.2	120.4	123.8	0.00	0.00 – 0.00	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	3.08	0.08 – 17.1	4
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	2.52	0.06 – 14.1	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	7.13	0.18 – 39.8	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	0.0	*	157.5	144.7	159.3	1.76	0.04 – 9.83	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	14.44	0.37 – 80.5	4

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Chagrin Falls Village

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	25.0	438.7	477.1	466.4	462.9	0.95	0.79 – 1.14	2
Bladder	2.0	29.2	21.5	21.6	21.0	1.54	0.74 – 2.83	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.88	0.39 – 5.50	4
Breast (Female)	3.8	122.7	122.2	120.4	123.8	0.99	0.60 – 1.55	2
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	2.6	37.0	51.1	52.1	49.1	0.87	0.47 – 1.50	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.10	0.30 – 2.82	1
Esophagus	<1	*	5.7	5.5	4.5	0.76	0.02 – 4.24	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.76	0.04 – 9.79	2
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.53	0.06 – 1.92	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	1.2	30.6	11.8	11.1	12.2	1.75	0.64 – 3.81	4
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	2.8	51.0	70.7	75.0	63.1	0.75	0.41 – 1.26	2
Melanoma of the Skin	1.4	28.1	15.3	17.6	19.6	1.35	0.54 – 2.78	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	1.6	26.7	19.1	19.2	19.5	1.44	0.62 – 2.84	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.71	0.09 – 2.55	2
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	1.12	0.30 – 2.86	4
Prostate (Male)	3.2	141.4	157.5	144.7	159.3	0.87	0.50 – 1.41	3
Stomach	<1	*	8.1	6.1	7.9	0.84	0.10 – 3.03	2
Testis (Male)	<1	*	4.9	5.5	5.4	2.25	0.06 – 12.5	4
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Cleveland Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	221.0	459.3	477.1	466.4	462.9	0.99	0.93 – 1.05	2
Bladder	6.0	12.6	21.5	21.6	21.0	0.60	0.41 – 0.86	1
Brain and Other CNS	2.6	5.4	5.8	6.8	6.4	0.82	0.44 – 1.41	1
Breast (Female)	39.8	141.4	122.2	120.4	123.8	1.14	0.98 – 1.30	4
Cervix (Female)	1.4	5.2	8.0	8.0	8.2	0.61	0.24 – 1.25	1
Colon & Rectum	23.8	50.6	51.1	52.1	49.1	1.02	0.84 – 1.21	1
Corpus Uterus (Female)	9.0	32.0	29.9	26.4	23.3	1.37	1.00 – 1.83	2
Esophagus	1.6	3.5	5.7	5.5	4.5	0.74	0.32 – 1.46	1
Hodgkin's Lymphoma	1.0	1.7	2.9	3.0	2.8	0.71	0.23 – 1.65	2
Kidney & Renal Pelvis	7.2	15.1	14.1	14.1	13.6	1.09	0.76 – 1.51	3
Larynx	1.8	3.7	4.7	4.4	3.5	1.07	0.49 – 2.03	1
Leukemia	5.6	12.0	11.8	11.1	12.2	0.96	0.64 – 1.39	3
Liver & Intrahepatic Bile Duct	1.6	3.3	5.6	4.3	6.6	0.49	0.21 – 0.97	1
Lung & Bronchus	25.2	53.1	70.7	75.0	63.1	0.84	0.70 – 1.00	1
Melanoma of the Skin	5.8	11.9	15.3	17.6	19.6	0.60	0.40 – 0.86	2
Multiple Myeloma	2.6	5.6	6.0	5.1	5.6	0.96	0.51 – 1.65	2
Non-Hodgkin's Lymphoma	8.6	17.9	19.1	19.2	19.5	0.91	0.66 – 1.23	2
Oral Cavity & Pharynx	3.0	6.1	10.4	9.6	10.4	0.59	0.33 – 0.97	2
Ovary	2.8	10	12.3	12.3	13.1	0.76	0.41 – 1.27	1
Pancreas	5.8	12.1	12.3	10.9	11.7	1.05	0.70 – 1.51	1
Prostate (Male)	41.4	204.6	157.5	144.7	159.3	1.32	1.14 – 1.51	3
Stomach	4.0	8.4	8.1	6.1	7.9	1.06	0.65 – 1.64	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.15	0.00 – 0.83	2
Thyroid	4.0	7.7	7.9	8.3	9.6	0.82	0.50 – 1.26	3

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Cuyahoga Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	4.0	427.8	477.1	466.4	462.9	0.82	0.49 – 1.30	1
Bladder	0.0	*	21.5	21.6	21.0	1.76	0.21 – 6.37	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	0.0	*	122.2	120.4	123.8	0.00	0.00 – 0.00	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	0.79	0.10 – 2.87	1
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	4.83	1.00 – 14.1	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	1.74	0.04 – 9.72	4
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.61	0.07 – 2.20	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	1.20	0.03 – 6.70	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	2.22	0.06 – 12.4	4
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	1.63	0.04 – 9.08	4
Prostate (Male)	0.0	*	157.5	144.7	159.3	1.18	0.32 – 3.03	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: East Cleveland

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	131.0	476.5	477.1	466.4	462.9	1.03	0.95 – 1.11	2
Bladder	3.0	12.1	21.5	21.6	21.0	0.59	0.34 – 0.94	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.34	0.07 – 0.99	2
Breast (Female)	18.0	112.4	122.2	120.4	123.8	0.89	0.72 – 1.10	3
Cervix (Female)	1.6	10.5	8.0	8.0	8.2	1.29	0.56 – 2.53	3
Colon & Rectum	14.6	53.9	51.1	52.1	49.1	1.08	0.85 – 1.36	3
Corpus Uterus (Female)	3.4	19.1	29.9	26.4	23.3	0.88	0.51 – 1.41	2
Esophagus	2.6	9.2	5.7	5.5	4.5	2.07	1.10 – 3.53	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.27	0.01 – 1.53	2
Kidney & Renal Pelvis	5.0	17.6	14.1	14.1	13.6	1.33	0.86 – 1.96	3
Larynx	1.4	5.3	4.7	4.4	3.5	1.43	0.57 – 2.95	2
Leukemia	2.2	8.2	11.8	11.1	12.2	0.65	0.33 – 1.17	2
Liver & Intrahepatic Bile Duct	2.2	8.0	5.6	4.3	6.6	1.21	0.60 – 2.16	2
Lung & Bronchus	22.2	81.0	70.7	75.0	63.1	1.25	1.03 – 1.51	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.15	0.04 – 0.39	2
Multiple Myeloma	3.2	11.8	6.0	5.1	5.6	2.06	1.18 – 3.34	4
Non-Hodgkin's Lymphoma	2.2	8.1	19.1	19.2	19.5	0.41	0.21 – 0.74	2
Oral Cavity & Pharynx	2.6	9.2	10.4	9.6	10.4	0.91	0.49 – 1.56	1
Ovary	2.0	12.3	12.3	12.3	13.1	0.93	0.45 – 1.71	4
Pancreas	3.6	13.0	12.3	10.9	11.7	1.12	0.66 – 1.77	3
Prostate (Male)	24.2	215.8	157.5	144.7	159.3	1.36	1.13 – 1.63	4
Stomach	4.0	14.5	8.1	6.1	7.9	1.85	1.13 – 2.85	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.34	0.01 – 1.92	3
Thyroid	1.0	3.7	7.9	8.3	9.6	0.40	0.13 – 0.93	2

Peer Comparison Neighborhoods/Municipalities: Central, Detroit-Shoreway, East Cleveland, Fairfax, Forest Hills, Glenville, Goodrich-Kirtland Park, Hough, Kinsman, North Broadway, Ohio City, St.Clair-Superior, Stockyards, Tremont, Union-Miles, University, Woodland Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Euclid

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	298.0	437.6	477.1	466.4	462.9	0.94	0.89 – 0.98	1
Bladder	14.0	18.4	21.5	21.6	21.0	0.87	0.67 – 1.10	2
Brain and Other CNS	3.2	5.6	5.8	6.8	6.4	0.81	0.46 – 1.31	3
Breast (Female)	43.4	116.7	122.2	120.4	123.8	0.94	0.82 – 1.08	3
Cervix (Female)	1.6	4.6	8.0	8.0	8.2	0.61	0.26 – 1.20	3
Colon & Rectum	38.4	53	51.1	52.1	49.1	1.07	0.92 – 1.23	1
Corpus Uterus (Female)	9.4	26.7	29.9	26.4	23.3	1.09	0.80 – 1.44	3
Esophagus	4.4	7.4	5.7	5.5	4.5	1.38	0.87 – 2.09	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.52	0.14 – 1.33	3
Kidney & Renal Pelvis	6.8	10.3	14.1	14.1	13.6	0.75	0.52 – 1.05	3
Larynx	1.8	2.6	4.7	4.4	3.5	0.78	0.36 – 1.49	1
Leukemia	6.0	9.0	11.8	11.1	12.2	0.71	0.48 – 1.01	3
Liver & Intrahepatic Bile Duct	4.6	6.7	5.6	4.3	6.6	1.04	0.66 – 1.56	3
Lung & Bronchus	48.6	68.3	70.7	75.0	63.1	1.06	0.93 – 1.20	1
Melanoma of the Skin	7.2	11.0	15.3	17.6	19.6	0.57	0.40 – 0.79	3
Multiple Myeloma	2.2	3.0	6.0	5.1	5.6	0.54	0.27 – 0.97	2
Non-Hodgkin's Lymphoma	11.6	16.2	19.1	19.2	19.5	0.86	0.65 – 1.11	3
Oral Cavity & Pharynx	6.8	11.9	10.4	9.6	10.4	1.02	0.70 – 1.42	3
Ovary	3.8	9.8	12.3	12.3	13.1	0.75	0.45 – 1.17	3
Pancreas	9.4	13.4	12.3	10.9	11.7	1.08	0.79 – 1.44	3
Prostate (Male)	36.6	138.5	157.5	144.7	159.3	0.85	0.73 – 0.98	3
Stomach	5.0	6.9	8.1	6.1	7.9	0.86	0.56 – 1.27	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.61	0.17 – 1.57	3
Thyroid	4.4	7.7	7.9	8.3	9.6	0.81	0.51 – 1.23	3

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Fairview Park

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	101.0	435.9	477.1	466.4	462.9	0.93	0.85 – 1.02	2
Bladder	4.0	17.7	21.5	21.6	21.0	0.79	0.49 – 1.21	1
Brain and Other CNS	1.2	4.5	5.8	6.8	6.4	0.89	0.33 – 1.94	2
Breast (Female)	11.0	94.1	122.2	120.4	123.8	0.71	0.54 – 0.92	1
Cervix (Female)	<1	*	8.0	8.0	8.2	0.68	0.14 – 2.00	3
Colon & Rectum	13.0	53.6	51.1	52.1	49.1	1.08	0.83 – 1.37	4
Corpus Uterus (Female)	4.6	36.5	29.9	26.4	23.3	1.57	0.99 – 2.35	3
Esophagus	1.2	5.3	5.7	5.5	4.5	1.10	0.40 – 2.40	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.58	0.43 – 4.04	3
Kidney & Renal Pelvis	3.0	12.1	14.1	14.1	13.6	0.96	0.54 – 1.58	3
Larynx	<1	*	4.7	4.4	3.5	0.74	0.15 – 2.18	3
Leukemia	2.4	10.2	11.8	11.1	12.2	0.85	0.44 – 1.49	2
Liver & Intrahepatic Bile Duct	1.2	5.1	5.6	4.3	6.6	0.78	0.29 – 1.71	4
Lung & Bronchus	14.4	58.9	70.7	75.0	63.1	0.92	0.72 – 1.16	4
Melanoma of the Skin	5	24.2	15.3	17.6	19.6	1.16	0.75 – 1.71	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.58	0.16 – 1.49	2
Non-Hodgkin's Lymphoma	5.6	23.6	19.1	19.2	19.5	1.22	0.81 – 1.77	3
Oral Cavity & Pharynx	1.6	7.4	10.4	9.6	10.4	0.69	0.30 – 1.36	1
Ovary	1.8	14.5	12.3	12.3	13.1	1.07	0.49 – 2.03	4
Pancreas	2.8	12.3	12.3	10.9	11.7	0.96	0.53 – 1.62	4
Prostate (Male)	13.4	137.4	157.5	144.7	159.3	0.84	0.65 – 1.07	2
Stomach	1.0	4.0	8.1	6.1	7.9	0.52	0.17 – 1.20	1
Testis (Male)	<1	*	4.9	5.5	5.4	1.40	0.29 – 4.09	3
Thyroid	1.8	8.8	7.9	8.3	9.6	0.96	0.44 – 1.82	3

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Garfield Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	172.0	452.3	477.1	466.4	462.9	0.95	0.89 – 1.02	1
Bladder	9.0	20.6	21.5	21.6	21.0	1.00	0.73 – 1.34	2
Brain and Other CNS	2.4	6.9	5.8	6.8	6.4	1.06	0.55 – 1.84	3
Breast (Female)	22.4	108.7	122.2	120.4	123.8	0.86	0.71 – 1.04	3
Cervix (Female)	1.8	11.3	8.0	8.0	8.2	1.22	0.56 – 2.32	4
Colon & Rectum	22.0	54.9	51.1	52.1	49.1	1.09	0.89 – 1.31	3
Corpus Uterus (Female)	5.4	25.2	29.9	26.4	23.3	1.11	0.73 – 1.61	1
Esophagus	2.6	6.9	5.7	5.5	4.5	1.44	0.77 – 2.46	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.45	0.05 – 1.63	1
Kidney & Renal Pelvis	7.0	19.7	14.1	14.1	13.6	1.36	0.94 – 1.89	4
Larynx	1.2	3.1	4.7	4.4	3.5	0.91	0.34 – 1.99	1
Leukemia	3.2	8.5	11.8	11.1	12.2	0.67	0.38 – 1.09	1
Liver & Intrahepatic Bile Duct	1.6	3.7	5.6	4.3	6.6	0.64	0.27 – 1.25	2
Lung & Bronchus	26.8	69.4	70.7	75.0	63.1	1.02	0.86 – 1.21	2
Melanoma of the Skin	3.8	10.3	15.3	17.6	19.6	0.53	0.32 – 0.83	1
Multiple Myeloma	2.2	5.6	6.0	5.1	5.6	0.95	0.48 – 1.71	2
Non-Hodgkin's Lymphoma	10.0	25.0	19.1	19.2	19.5	1.30	0.97 – 1.72	4
Oral Cavity & Pharynx	3.8	11.8	10.4	9.6	10.4	1.00	0.60 – 1.56	3
Ovary	2.0	10.8	12.3	12.3	13.1	0.70	0.34 – 1.28	2
Pancreas	4.6	12.2	12.3	10.9	11.7	0.94	0.60 – 1.41	2
Prostate (Male)	18.4	123.1	157.5	144.7	159.3	0.74	0.60 – 0.91	2
Stomach	3.8	10.0	8.1	6.1	7.9	1.16	0.70 – 1.82	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.06	0.29 – 2.71	4
Thyroid	2.4	7.6	7.9	8.3	9.6	0.77	0.40 – 1.35	3

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Gates Mills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	12.0	375.2	477.1	466.4	462.9	0.75	0.58 – 0.97	1
Bladder	0.0	*	21.5	21.6	21.0	0.00	0.00 – 0.00	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.6	89.7	122.2	120.4	123.8	0.73	0.32 – 1.44	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.4	46.4	51.1	52.1	49.1	0.80	0.32 – 1.65	2
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.43	0.29 – 4.17	2
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.81	0.1 – 2.94	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.50	0.01 – 2.81	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.82	0.02 – 4.58	2
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.26	0.05 – 0.75	1
Melanoma of the Skin	1.8	51.2	15.3	17.6	19.6	2.77	1.27 – 5.26	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.98	0.02 – 5.46	2
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.59	0.07 – 2.15	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	2.8	141.6	157.5	144.7	159.3	0.90	0.49 – 1.52	1
Stomach	<1	*	8.1	6.1	7.9	1.44	0.17 – 5.20	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	1.41	0.17 – 5.11	4

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Glenwillow

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	7.0	1599.6	477.1	466.4	462.9	3.06	2.14 – 4.24	4
Bladder	1.0	113.8	21.5	21.6	21.0	5.78	1.2 – 16.89	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.0	500.0	122.2	120.4	123.8	3.22	1.05 – 7.52	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	3.24	0.88 – 8.30	4
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	<1	*	5.7	5.5	4.5	8.68	0.2 – 48.38	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	16.04	0.41 – 89.4	4
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	2.86	0.07 – 15.9	4
Larynx	<1	*	4.7	4.4	3.5	10.95	0.30 – 61.0	4
Leukemia	<1	*	11.8	11.1	12.2	3.41	0.1 – 18.99	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	5.90	0.15 – 32.9	4
Lung & Bronchus	1.0	231.3	70.7	75.0	63.1	3.18	1.03 – 7.43	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	2.01	0.05 – 11.2	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	2.04	0.05 – 11.4	4
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	3.42	0.09 – 19.1	4
Prostate (Male)	0.0	405.3	157.5	144.7	159.3	2.39	0.77 – 5.57	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Highland Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	56.0	556.2	477.1	466.4	462.9	1.15	1.02 – 1.29	4
Bladder	4.0	35.8	21.5	21.6	21.0	1.75	1.07 – 2.70	4
Brain and Other CNS	1.6	15.4	5.8	6.8	6.4	2.61	1.13 – 5.15	4
Breast (Female)	6.2	121.4	122.2	120.4	123.8	0.93	0.63 – 1.32	1
Cervix (Female)	<1	*	8.0	8.0	8.2	0.53	0.01 – 2.94	3
Colon & Rectum	5.0	47.2	51.1	52.1	49.1	0.95	0.62 – 1.41	2
Corpus Uterus (Female)	2.0	38.2	29.9	26.4	23.3	1.59	0.76 – 2.92	4
Esophagus	<1	*	5.7	5.5	4.5	1.65	0.45 – 4.21	4
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	2.2	21.8	14.1	14.1	13.6	1.54	0.77 – 2.76	4
Larynx	<1	*	4.7	4.4	3.5	0.54	0.01 – 3.01	2
Leukemia	1.8	18.1	11.8	11.1	12.2	1.47	0.67 – 2.78	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.14	0.31 – 2.93	4
Lung & Bronchus	6.8	64.0	70.7	75.0	63.1	0.98	0.68 – 1.37	4
Melanoma of the Skin	2.6	33.1	15.3	17.6	19.6	1.35	0.72 – 2.30	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.32	0.36 – 3.37	4
Non-Hodgkin's Lymphoma	3.2	30.1	19.1	19.2	19.5	1.58	0.90 – 2.57	4
Oral Cavity & Pharynx	1.0	9.4	10.4	9.6	10.4	0.94	0.30 – 2.19	3
Ovary	1.2	23.8	12.3	12.3	13.1	1.69	0.62 – 3.69	4
Pancreas	2.2	22.2	12.3	10.9	11.7	1.75	0.87 – 3.13	4
Prostate (Male)	7.4	150.5	157.5	144.7	159.3	0.91	0.64 – 1.26	3
Stomach	<1	*	8.1	6.1	7.9	0.48	0.06 – 1.72	1
Testis (Male)	<1	*	4.9	5.5	5.4	2.34	0.28 – 8.47	4
Thyroid	1.4	14.4	7.9	8.3	9.6	1.65	0.66 – 3.40	4

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Highland Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	9.0	480.8	477.1	466.4	462.9	1.02	0.74 – 1.37	2
Bladder	0.0	*	21.5	21.6	21.0	0.00	0.00 – 0.00	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	<1	*	122.2	120.4	123.8	0.74	0.20 – 1.88	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.4	74.6	51.1	52.1	49.1	1.52	0.61 – 3.13	3
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.95	0.02 – 5.31	2
Esophagus	<1	*	5.7	5.5	4.5	2.39	0.06 – 13.3	4
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.92	0.02 – 5.11	4
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	1.4	76.9	70.7	75.0	63.1	1.18	0.48 – 2.44	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.58	0.01 – 3.21	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	1.12	0.14 – 4.04	4
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	<1	*	12.3	12.3	13.1	1.69	0.04 – 9.41	4
Pancreas	<1	*	12.3	10.9	11.7	1.82	0.22 – 6.56	3
Prostate (Male)	1.2	154.6	157.5	144.7	159.3	0.88	0.32 – 1.92	3
Stomach	<1	*	8.1	6.1	7.9	2.71	0.33 – 9.79	4
Testis (Male)	<1	*	4.9	5.5	5.4	3.25	0.08 – 18.1	4
Thyroid	<1	*	7.9	8.3	9.6	1.25	0.03 – 6.94	4

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Hunting Valley

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	5.0	494.0	477.1	466.4	462.9	1.02	0.65 – 1.52	2
Bladder	0.0	*	21.5	21.6	21.0	1.79	0.22 – 6.45	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.4	376.0	122.2	120.4	123.8	2.32	0.93 – 4.78	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	0.79	0.10 – 2.84	1
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 – 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.87	0.18 – 2.55	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	4.51	1.23 – 11.6	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 – 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	3.25	0.39 – 11.7	4
Prostate (Male)	0.0	*	157.5	144.7	159.3	0.89	0.24 – 2.28	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Independence

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	46.0	449.8	477.1	466.4	462.9	0.97	0.85 – 1.11	1
Bladder	2.0	15.8	21.5	21.6	21.0	0.76	0.35 – 1.45	2
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.69	0.08 – 2.51	1
Breast (Female)	7.0	119.3	122.2	120.4	123.8	1.07	0.75 – 1.49	1
Cervix (Female)	<1	*	8.0	8.0	8.2	0.57	0.01 – 3.20	3
Colon & Rectum	5.2	52.8	51.1	52.1	49.1	0.98	0.64 – 1.43	4
Corpus Uterus (Female)	1.4	25.0	29.9	26.4	23.3	1.14	0.46 – 2.34	1
Esophagus	<1	*	5.7	5.5	4.5	0.83	0.10 – 3.00	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.98	0.24 – 7.16	4
Kidney & Renal Pelvis	1.4	14.1	14.1	14.1	13.6	1.02	0.41 – 2.1	3
Larynx	<1	*	4.7	4.4	3.5	0.56	0.01 – 3.13	3
Leukemia	1	10.5	11.8	11.1	12.2	0.82	0.27 – 1.91	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.30	0.01 – 1.65	1
Lung & Bronchus	6.0	50.2	70.7	75.0	63.1	0.86	0.58 – 1.23	2
Melanoma of the Skin	2.2	21.4	15.3	17.6	19.6	1.19	0.59 – 2.12	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.98	0.20 – 2.87	3
Non-Hodgkin's Lymphoma	2.6	24.6	19.1	19.2	19.5	1.30	0.69 – 2.22	3
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.59	0.12 – 1.73	2
Ovary	1.0	20.1	12.3	12.3	13.1	1.41	0.46 – 3.28	4
Pancreas	2.0	17.5	12.3	10.9	11.7	1.56	0.75 – 2.86	3
Prostate (Male)	6.2	146.7	157.5	144.7	159.3	0.84	0.57 – 1.20	2
Stomach	1.0	7.7	8.1	6.1	7.9	1.17	0.38 – 2.73	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	1.0	13.0	7.9	8.3	9.6	1.30	0.42 – 3.04	4

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Lakewood

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	244.0	455.1	477.1	466.4	462.9	0.98	0.93 – 1.04	2
Bladder	13.0	24.4	21.5	21.6	21.0	1.11	0.86 – 1.42	3
Brain and Other CNS	3.2	6.1	5.8	6.8	6.4	0.91	0.52 – 1.48	2
Breast (Female)	36.4	121.6	122.2	120.4	123.8	0.97	0.83 – 1.12	3
Cervix (Female)	2.6	9.0	8.0	8.0	8.2	1.03	0.55 – 1.76	3
Colon & Rectum	27.8	50.0	51.1	52.1	49.1	1.03	0.87 – 1.22	1
Corpus Uterus (Female)	10.6	37.6	29.9	26.4	23.3	1.53	1.14 – 2.00	3
Esophagus	3.0	5.7	5.7	5.5	4.5	1.25	0.70 – 2.07	2
Hodgkin's Lymphoma	1.4	2.3	2.9	3.0	2.8	0.84	0.34 – 1.72	2
Kidney & Renal Pelvis	6.8	12.7	14.1	14.1	13.6	0.95	0.65 – 1.32	2
Larynx	2.2	4.4	4.7	4.4	3.5	1.22	0.61 – 2.19	2
Leukemia	9.6	17	11.8	11.1	12.2	1.43	1.06 – 1.90	4
Liver & Intrahepatic Bile Duct	2.2	4.2	5.6	4.3	6.6	0.63	0.31 – 1.12	3
Lung & Bronchus	31.4	59.5	70.7	75.0	63.1	0.94	0.80 – 1.09	1
Melanoma of the Skin	9.8	17.4	15.3	17.6	19.6	0.90	0.67 – 1.19	3
Multiple Myeloma	3.0	5.8	6.0	5.1	5.6	0.99	0.55 – 1.63	3
Non-Hodgkin's Lymphoma	8.8	15.4	19.1	19.2	19.5	0.82	0.60 – 1.10	1
Oral Cavity & Pharynx	7.0	13.5	10.4	9.6	10.4	1.27	0.88 – 1.76	4
Ovary	3.6	12.1	12.3	12.3	13.1	0.88	0.52 – 1.39	3
Pancreas	5.6	10.2	12.3	10.9	11.7	0.88	0.58 – 1.27	1
Prostate (Male)	23.6	115.3	157.5	144.7	159.3	0.73	0.60 – 0.87	1
Stomach	3.4	6.2	8.1	6.1	7.9	0.78	0.45 – 1.25	2
Testis (Male)	2.8	8.1	4.9	5.5	5.4	1.61	0.88 – 2.70	4
Thyroid	4.4	7.3	7.9	8.3	9.6	0.79	0.50 – 1.20	2

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Linndale

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	1.0	1504.0	477.1	466.4	462.9	2.20	0.72 — 5.15	4
Bladder	0.0	*	21.5	21.6	21.0	10.93	0.28 — 60.9	4
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 — 0.00	1
Breast (Female)	<1	*	122.2	120.4	123.8	2.7	0.07 — 15.1	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 — 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	4.6	0.12 — 25.6	4
Corpus Uterus (Female)	0.0	*	29.9	26.4	23.3	0.00	0.00 — 0.00	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 — 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 — 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 — 0.00	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 — 0.00	1
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 — 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 — 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.00	0.00 — 0.00	1
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 — 0.00	1
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 — 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 — 0.00	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 — 0.00	1
Ovary	0.0	*	12.3	12.3	13.1	0.00	0.00 — 0.00	1
Pancreas	<1	*	12.3	10.9	11.7	19.76	0.5 — 110.1	4
Prostate (Male)	0.0	*	157.5	144.7	159.3	3.06	0.08 — 17.0	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 — 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 — 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 — 0.00	1

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Lyndhurst

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	108.0	420.6	477.1	466.4	462.9	0.91	0.83 – 0.99	1
Bladder	6.0	21.1	21.5	21.6	21.0	0.93	0.62 – 1.34	2
Brain and Other CNS	1.8	9.2	5.8	6.8	6.4	1.32	0.61 – 2.51	4
Breast (Female)	20.0	142.1	122.2	120.4	123.8	1.25	1.02 – 1.52	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.49	0.06 – 1.79	2
Colon & Rectum	12.4	45.5	51.1	52.1	49.1	0.90	0.69 – 1.15	1
Corpus Uterus (Female)	4.2	36.7	29.9	26.4	23.3	1.37	0.85 – 2.10	4
Esophagus	2.0	6.8	5.7	5.5	4.5	1.65	0.79 – 3.03	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.75	0.48 – 4.47	4
Kidney & Renal Pelvis	3.8	15.3	14.1	14.1	13.6	1.13	0.68 – 1.77	3
Larynx	<1	*	4.7	4.4	3.5	0.93	0.25 – 2.39	4
Leukemia	2.0	5.4	11.8	11.1	12.2	0.64	0.31 – 1.18	1
Liver & Intrahepatic Bile Duct	1.2	4.2	5.6	4.3	6.6	0.73	0.27 – 1.59	1
Lung & Bronchus	11.6	41.4	70.7	75.0	63.1	0.65	0.50 – 0.84	1
Melanoma of the Skin	4.2	16.5	15.3	17.6	19.6	0.94	0.58 – 1.43	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.38	0.08 – 1.12	1
Non-Hodgkin's Lymphoma	4.8	17.8	19.1	19.2	19.5	0.95	0.61 – 1.42	2
Oral Cavity & Pharynx	1.8	6.8	10.4	9.6	10.4	0.74	0.34 – 1.41	1
Ovary	1.6	9.3	12.3	12.3	13.1	0.89	0.39 – 1.76	2
Pancreas	2.8	10.7	12.3	10.9	11.7	0.83	0.45 – 1.39	1
Prostate (Male)	13.6	121.0	157.5	144.7	159.3	0.78	0.60 – 0.98	1
Stomach	2.6	8.7	8.1	6.1	7.9	1.17	0.62 – 2.00	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.19	0.14 – 4.31	4
Thyroid	1.6	10.1	7.9	8.3	9.6	0.94	0.41 – 1.85	2

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Maple Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	141.0	478.4	477.1	466.4	462.9	1.01	0.94 – 1.09	4
Bladder	8.0	25.6	21.5	21.6	21.0	1.21	0.87 – 1.65	3
Brain and Other CNS	1.8	5.8	5.8	6.8	6.4	0.98	0.45 – 1.87	3
Breast (Female)	18.8	116.1	122.2	120.4	123.8	0.92	0.74 – 1.13	2
Cervix (Female)	2.0	15.0	8.0	8.0	8.2	1.62	0.77 – 2.97	4
Colon & Rectum	16.6	56.6	51.1	52.1	49.1	1.08	0.86 – 1.34	4
Corpus Uterus (Female)	5.2	32.7	29.9	26.4	23.3	1.36	0.89 – 1.99	3
Esophagus	2.0	6.5	5.7	5.5	4.5	1.44	0.69 – 2.65	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.54	0.07 – 1.97	2
Kidney & Renal Pelvis	5.0	16.6	14.1	14.1	13.6	1.24	0.8 – 1.83	3
Larynx	1.2	4.1	4.7	4.4	3.5	1.17	0.43 – 2.54	3
Leukemia	4.0	13.3	11.8	11.1	12.2	1.08	0.66 – 1.67	3
Liver & Intrahepatic Bile Duct	2.0	7.3	5.6	4.3	6.6	1.02	0.49 – 1.87	4
Lung & Bronchus	18.4	60.2	70.7	75.0	63.1	0.92	0.74 – 1.13	2
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.14	0.04 – 0.36	1
Multiple Myeloma	2.4	7.4	6.0	5.1	5.6	1.36	0.70 – 2.38	4
Non-Hodgkin's Lymphoma	4.6	15.2	19.1	19.2	19.5	0.77	0.49 – 1.16	2
Oral Cavity & Pharynx	2.2	8.1	10.4	9.6	10.4	0.73	0.37 – 1.31	2
Ovary	3.2	18.5	12.3	12.3	13.1	1.45	0.83 – 2.35	4
Pancreas	3.0	10.6	12.3	10.9	11.7	0.81	0.45 – 1.34	2
Prostate (Male)	23	192.8	157.5	144.7	159.3	1.16	0.96 – 1.40	4
Stomach	3.0	9.9	8.1	6.1	7.9	1.21	0.68 – 2.00	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.62	0.08 – 2.24	2
Thyroid	1.8	7.1	7.9	8.3	9.6	0.69	0.32 – 1.31	3

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Mayfield Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	141.0	438.8	477.1	466.4	462.9	0.92	0.85 – 0.99	2
Bladder	9.0	27.5	21.5	21.6	21.0	1.15	0.85 – 1.53	3
Brain and Other CNS	2.2	9.4	5.8	6.8	6.4	1.27	0.63 – 2.27	4
Breast (Female)	21.6	130.8	122.2	120.4	123.8	1.02	0.84 – 1.23	4
Cervix (Female)	1.0	9.2	8.0	8.0	8.2	0.96	0.31 – 2.24	4
Colon & Rectum	13.4	37.2	51.1	52.1	49.1	0.74	0.57 – 0.94	1
Corpus Uterus (Female)	5.6	33.7	29.9	26.4	23.3	1.38	0.92 – 2.00	4
Esophagus	1.4	3.1	5.7	5.5	4.5	0.89	0.36 – 1.83	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.65	0.08 – 2.35	2
Kidney & Renal Pelvis	4.4	17.2	14.1	14.1	13.6	1.02	0.64 – 1.55	3
Larynx	<1	*	4.7	4.4	3.5	0.54	0.11 – 1.59	2
Leukemia	4.0	13.6	11.8	11.1	12.2	0.99	0.60 – 1.52	3
Liver & Intrahepatic Bile Duct	1.0	2.8	5.6	4.3	6.6	0.48	0.15 – 1.11	2
Lung & Bronchus	21.0	58.1	70.7	75.0	63.1	0.91	0.74 – 1.10	2
Melanoma of the Skin	4.6	14.2	15.3	17.6	19.6	0.80	0.50 – 1.20	2
Multiple Myeloma	2.2	6.1	6.0	5.1	5.6	1.08	0.54 – 1.93	2
Non-Hodgkin's Lymphoma	6.2	18.1	19.1	19.2	19.5	0.94	0.64 – 1.34	3
Oral Cavity & Pharynx	2.8	8.8	10.4	9.6	10.4	0.90	0.49 – 1.51	2
Ovary	2.0	10.7	12.3	12.3	13.1	0.83	0.40 – 1.52	3
Pancreas	5.8	17.1	12.3	10.9	11.7	1.31	0.88 – 1.88	4
Prostate (Male)	13.4	102.3	157.5	144.7	159.3	0.63	0.49 – 0.80	1
Stomach	3.6	9.9	8.1	6.1	7.9	1.24	0.73 – 1.96	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.67	0.45 – 4.27	4
Thyroid	1.8	5.7	7.9	8.3	9.6	0.83	0.38 – 1.57	2

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Mayfield Village

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	22.0	408.4	477.1	466.4	462.9	0.87	0.71 – 1.05	1
Bladder	1.0	18.4	21.5	21.6	21.0	0.82	0.26 – 1.90	2
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	3.6	142.1	122.2	120.4	123.8	1.10	0.65 – 1.73	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	2.6	46.3	51.1	52.1	49.1	0.94	0.50 – 1.61	2
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.26	0.34 – 3.23	2
Esophagus	<1	*	5.7	5.5	4.5	0.78	0.02 – 4.35	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.82	0.17 – 2.41	2
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.64	0.08 – 2.32	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.56	0.01 – 3.13	4
Lung & Bronchus	2.8	47.9	70.7	75.0	63.1	0.76	0.41 – 1.27	1
Melanoma of the Skin	1.2	22	15.3	17.6	19.6	1.25	0.46 – 2.73	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.62	0.02 – 3.48	1
Non-Hodgkin's Lymphoma	1.8	41.6	19.1	19.2	19.5	1.73	0.79 – 3.29	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.37	0.01 – 2.08	1
Ovary	<1	*	12.3	12.3	13.1	1.13	0.14 – 4.08	4
Pancreas	<1	*	12.3	10.9	11.7	0.60	0.07 – 2.16	1
Prostate (Male)	3.0	119.1	157.5	144.7	159.3	0.68	0.38 – 1.13	1
Stomach	<1	*	8.1	6.1	7.9	0.45	0.01 – 2.52	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.51	0.01 – 2.86	1

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Middleburg Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	94.0	414.1	477.1	466.4	462.9	0.88	0.80 – 0.96	1
Bladder	5.0	22.4	21.5	21.6	21.0	1.03	0.68 – 1.50	3
Brain and Other CNS	1.8	9.1	5.8	6.8	6.4	1.41	0.64 – 2.67	4
Breast (Female)	12.6	111.1	122.2	120.4	123.8	0.85	0.66 – 1.09	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	11.6	49.2	51.1	52.1	49.1	0.98	0.74 – 1.27	3
Corpus Uterus (Female)	3.6	31.6	29.9	26.4	23.3	1.26	0.74 – 1.98	3
Esophagus	1.4	5.9	5.7	5.5	4.5	1.29	0.52 – 2.65	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.42	0.01 – 2.37	1
Kidney & Renal Pelvis	3.4	15.3	14.1	14.1	13.6	1.10	0.64 – 1.76	4
Larynx	<1	*	4.7	4.4	3.5	0.74	0.15 – 2.15	3
Leukemia	2.0	8.2	11.8	11.1	12.2	0.74	0.36 – 1.37	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.26	0.03 – 0.96	2
Lung & Bronchus	15.0	60.1	70.7	75.0	63.1	0.96	0.76 – 1.20	4
Melanoma of the Skin	4.2	21.1	15.3	17.6	19.6	1.01	0.62 – 1.54	3
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	3.6	16.5	19.1	19.2	19.5	0.81	0.48 – 1.27	1
Oral Cavity & Pharynx	1.0	4.2	10.4	9.6	10.4	0.44	0.14 – 1.02	1
Ovary	1.4	11.7	12.3	12.3	13.1	0.87	0.35 – 1.79	3
Pancreas	1.6	6.5	12.3	10.9	11.7	0.56	0.24 – 1.10	1
Prostate (Male)	14.2	138.9	157.5	144.7	159.3	0.83	0.65 – 1.04	2
Stomach	1.2	5.1	8.1	6.1	7.9	0.63	0.23 – 1.38	1
Testis (Male)	<1	*	4.9	5.5	5.4	1.55	0.32 – 4.53	4
Thyroid	1.8	10.5	7.9	8.3	9.6	1.02	0.47 – 1.94	4

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Moreland Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	23.0	502.1	477.1	466.4	462.9	1.08	0.90 – 1.30	3
Bladder	2.0	31.4	21.5	21.6	21.0	1.64	0.71 – 3.22	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.75	0.02 – 4.17	2
Breast (Female)	3.4	159.5	122.2	120.4	123.8	1.12	0.65 – 1.80	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.4	28.3	51.1	52.1	49.1	0.61	0.25 – 1.26	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.68	0.08 – 2.47	1
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	1.4	27.9	14.1	14.1	13.6	2.16	0.87 – 4.45	4
Larynx	<1	*	4.7	4.4	3.5	2.31	0.28 – 8.34	4
Leukemia	<1	*	11.8	11.1	12.2	1.54	0.42 – 3.95	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.62	0.02 – 3.43	4
Lung & Bronchus	2.8	60.1	70.7	75.0	63.1	0.92	0.50 – 1.54	4
Melanoma of the Skin	2.8	62.6	15.3	17.6	19.6	3.25	1.77 – 5.45	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	2.24	0.46 – 6.56	4
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.68	0.14 – 1.98	2
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.80	0.10 – 2.90	3
Ovary	<1	*	12.3	12.3	13.1	1.27	0.15 – 4.59	4
Pancreas	0.0	*	12.3	10.9	11.7	0.00	0.00 – 0.00	1
Prostate (Male)	3.8	171.0	157.5	144.7	159.3	1.00	0.60 – 1.55	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.53	0.01 – 2.93	1

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Newburgh Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	13.0	502.6	477.1	466.4	462.9	1.07	0.82 – 1.37	3
Bladder	1.0	31.6	21.5	21.6	21.0	1.47	0.40 – 3.77	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	2.51	0.30 – 9.06	4
Breast (Female)	<1	*	122.2	120.4	123.8	0.34	0.07 – 0.98	1
Cervix (Female)	<1	*	8.0	8.0	8.2	1.83	0.1 – 10.19	4
Colon & Rectum	1.0	37.5	51.1	52.1	49.1	0.79	0.26 – 1.85	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.78	0.37 – 5.21	4
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	1.16	0.14 – 4.21	3
Larynx	<1	*	4.7	4.4	3.5	2.29	0.06 – 12.8	4
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	3.6	141.9	70.7	75.0	63.1	2.23	1.32 – 3.52	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.80	0.10 – 2.90	4
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.40	0.01 – 2.24	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	1.53	0.19 – 5.53	4
Ovary	<1	*	12.3	12.3	13.1	1.04	0.03 – 5.78	4
Pancreas	<1	*	12.3	10.9	11.7	1.33	0.16 – 4.81	4
Prostate (Male)	2.4	232.7	157.5	144.7	159.3	1.55	0.80 – 2.71	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: North Olmsted

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	185.0	463.0	477.1	466.4	462.9	1.00	0.94 – 1.07	2
Bladder	9.0	22.5	21.5	21.6	21.0	1.09	0.80– 1.46	3
Brain and Other CNS	2.6	6.8	5.8	6.8	6.4	1.08	0.57 – 1.84	3
Breast (Female)	27.6	125.9	122.2	120.4	123.8	1.02	0.86 – 1.20	3
Cervix (Female)	<1	*	8.0	8.0	8.2	0.37	0.08 – 1.08	1
Colon & Rectum	20.0	50.3	51.1	52.1	49.1	1.02	0.83 – 1.24	3
Corpus Uterus (Female)	5.8	26.1	29.9	26.4	23.3	1.12	0.75 – 1.61	2
Esophagus	2.2	5.5	5.7	5.5	4.5	1.21	0.60 – 2.16	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.84	0.23 – 2.15	3
Kidney & Renal Pelvis	4.6	11.3	14.1	14.1	13.6	0.85	0.54 – 1.27	1
Larynx	2.0	4.9	4.7	4.4	3.5	1.41	0.68 – 2.60	4
Leukemia	5.8	15.5	11.8	11.1	12.2	1.24	0.83 – 1.79	4
Liver & Intrahepatic Bile Duct	2.2	5.4	5.6	4.3	6.6	0.82	0.41 – 1.48	3
Lung & Bronchus	24.4	59.7	70.7	75.0	63.1	0.96	0.79 – 1.14	3
Melanoma of the Skin	7.0	17.6	15.3	17.6	19.6	0.92	0.64 – 1.28	1
Multiple Myeloma	1.6	3.9	6.0	5.1	5.6	0.71	0.31 – 1.39	1
Non-Hodgkin's Lymphoma	9.2	23.2	19.1	19.2	19.5	1.20	0.88 – 1.60	3
Oral Cavity & Pharynx	2.8	7.3	10.4	9.6	10.4	0.68	0.37 – 1.13	1
Ovary	1.8	8.3	12.3	12.3	13.1	0.63	0.29 – 1.19	1
Pancreas	5.2	13	12.3	10.9	11.7	1.11	0.73 – 1.63	2
Prostate (Male)	27.2	156.3	157.5	144.7	159.3	0.96	0.81 – 1.14	3
Stomach	2.0	5.1	8.1	6.1	7.9	0.64	0.30 – 1.17	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.94	0.26 – 2.40	3
Thyroid	3.0	7.6	7.9	8.3	9.6	0.84	0.47 – 1.39	1

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: North Randall

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	10.0	881.9	477.1	466.4	462.9	1.98	1.48 – 2.60	4
Bladder	0.0	*	21.5	21.6	21.0	0.00	0.00 – 0.00	1
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	1.0	185.9	122.2	120.4	123.8	1.42	0.46 – 3.32	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.6	111.6	51.1	52.1	49.1	2.73	1.18 – 5.38	4
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	4.56	0.94 – 13.3	4
Esophagus	<1	*	5.7	5.5	4.5	3.85	0.1 – 21.47	4
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	0.0	*	14.1	14.1	13.6	0.00	0.00 – 0.00	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	2.2	188.2	70.7	75.0	63.1	3.01	1.50 – 5.38	4
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	3.03	0.08 – 16.8	4
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	5.25	1.08 – 15.3	4
Ovary	<1	*	12.3	12.3	13.1	2.66	0.07 – 14.8	4
Pancreas	<1	*	12.3	10.9	11.7	2.85	0.34 – 10.3	4
Prostate (Male)	2.0	345.7	157.5	144.7	159.3	2.29	1.10 – 4.20	4
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: North Royalton

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	145.0	509.1	477.1	466.4	462.9	1.09	1.01 – 1.17	3
Bladder	8.0	28.6	21.5	21.6	21.0	1.33	0.94 – 1.81	4
Brain and Other CNS	2.2	7.9	5.8	6.8	6.4	1.19	0.59 – 2.12	3
Breast (Female)	20.2	125.4	122.2	120.4	123.8	1.01	0.82 – 1.23	2
Cervix (Female)	1.6	10.2	8.0	8.0	8.2	1.22	0.53 – 2.40	4
Colon & Rectum	12.8	45.1	51.1	52.1	49.1	0.92	0.71 – 1.17	1
Corpus Uterus (Female)	6.2	38.8	29.9	26.4	23.3	1.67	1.14 – 2.37	4
Esophagus	1.6	5.9	5.7	5.5	4.5	1.24	0.54 – 2.45	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.25	0.01 – 1.41	2
Kidney & Renal Pelvis	4.0	14.0	14.1	14.1	13.6	1.01	0.62 – 1.56	2
Larynx	<1	*	4.7	4.4	3.5	0.59	0.12 – 1.74	2
Leukemia	5.0	18.1	11.8	11.1	12.2	1.46	0.95 – 2.16	4
Liver & Intrahepatic Bile Duct	2.0	7.2	5.6	4.3	6.6	1.03	0.49 – 1.90	4
Lung & Bronchus	18.0	65.4	70.7	75.0	63.1	1.01	0.81 – 1.24	4
Melanoma of the Skin	5.6	19.1	15.3	17.6	19.6	0.97	0.64 – 1.40	2
Multiple Myeloma	1.4	5.0	6.0	5.1	5.6	0.87	0.35 – 1.80	3
Non-Hodgkin's Lymphoma	6.0	21.4	19.1	19.2	19.5	1.07	0.72 – 1.53	2
Oral Cavity & Pharynx	2.8	9.3	10.4	9.6	10.4	0.91	0.50 – 1.53	3
Ovary	2.6	16.0	12.3	12.3	13.1	1.24	0.66 – 2.12	3
Pancreas	4.0	15.2	12.3	10.9	11.7	1.22	0.74 – 1.88	3
Prostate (Male)	20.6	171	157.5	144.7	159.3	1.04	0.85 – 1.26	4
Stomach	2.0	7.3	8.1	6.1	7.9	0.89	0.43 – 1.64	2
Testis (Male)	<1	*	4.9	5.5	5.4	0.26	0.01 – 1.45	2
Thyroid	3.4	11.3	7.9	8.3	9.6	1.17	0.68 – 1.87	3

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Oakwood

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	23.0	424.3	477.1	466.4	462.9	0.93	0.77 – 1.11	1
Bladder	1.0	13.7	21.5	21.6	21.0	0.65	0.18 – 1.67	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.68	0.02 – 3.78	1
Breast (Female)	2.2	75.4	122.2	120.4	123.8	0.63	0.32 – 1.13	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	3.6	58.0	51.1	52.1	49.1	1.29	0.76 – 2.03	4
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	0.30	0.01 – 1.69	1
Esophagus	<1	*	5.7	5.5	4.5	0.80	0.02 – 4.48	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.88	0.05 – 10.48	4
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	1.14	0.31 – 2.91	3
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.62	0.08 – 2.26	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.72	0.35 – 5.02	4
Lung & Bronchus	4.6	85.6	70.7	75.0	63.1	1.30	0.83 – 1.95	4
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.42	0.05 – 1.51	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.63	0.02 – 3.53	2
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	0.58	0.12 – 1.69	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	<1	*	12.3	12.3	13.1	0.52	0.01 – 2.90	2
Pancreas	<1	*	12.3	10.9	11.7	0.89	0.18 – 2.59	3
Prostate (Male)	4.8	215.2	157.5	144.7	159.3	1.35	0.86 – 2.00	4
Stomach	1.0	15.0	8.1	6.1	7.9	2.23	0.72 – 5.21	4
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0	*	7.9	8.3	9.6	0.0	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Olmsted Falls

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	43.0	599.7	477.1	466.4	462.9	1.29	1.12 – 1.47	4
Bladder	1.0	13.9	21.5	21.6	21.0	0.72	0.23 – 1.67	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.82	0.10 – 2.97	1
Breast (Female)	8.6	208.8	122.2	120.4	123.8	1.63	1.18 – 2.20	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	4.2	64.6	51.1	52.1	49.1	1.25	0.77 – 1.91	4
Corpus Uterus (Female)	1.6	34.2	29.9	26.4	23.3	1.65	0.71 – 3.25	3
Esophagus	<1	*	5.7	5.5	4.5	1.27	0.15 – 4.59	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	1.0	13.2	14.1	14.1	13.6	1.01	0.33 – 2.35	2
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.94	0.25 – 2.39	3
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.82	0.10 – 2.97	4
Lung & Bronchus	4.6	67.7	70.7	75.0	63.1	1.06	0.67 – 1.60	4
Melanoma of the Skin	2.4	31.0	15.3	17.6	19.6	1.62	0.84 – 2.83	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	2.05	0.56 – 5.25	4
Non-Hodgkin's Lymphoma	2.8	40.6	19.1	19.2	19.5	2.01	1.10 – 3.37	4
Oral Cavity & Pharynx	1.0	11.8	10.4	9.6	10.4	1.29	0.42 – 3.00	4
Ovary	1.2	32.1	12.3	12.3	13.1	2.22	0.81 – 4.83	4
Pancreas	1.4	17.9	12.3	10.9	11.7	1.79	0.72 – 3.68	4
Prostate (Male)	4.4	148.4	157.5	144.7	159.3	0.93	0.58 – 1.41	2
Stomach	<1	*	8.1	6.1	7.9	1.47	0.40 – 3.77	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.94	0.02 – 5.26	3
Thyroid	<1	*	7.9	8.3	9.6	0.76	0.16 – 2.22	1

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Olmsted Township

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	75.0	505.2	477.1	466.4	462.9	1.10	0.99 – 1.21	4
Bladder	3.0	16.7	21.5	21.6	21.0	0.77	0.41 – 1.32	1
Brain and Other CNS	1.2	12.3	5.8	6.8	6.4	1.44	0.53 – 3.13	4
Breast (Female)	11.0	133.5	122.2	120.4	123.8	1.13	0.85 – 1.48	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	9.0	54.3	51.1	52.1	49.1	1.18	0.86 – 1.57	4
Corpus Uterus (Female)	4.2	52.9	29.9	26.4	23.3	2.27	1.41 – 3.47	4
Esophagus	<1	*	5.7	5.5	4.5	1.16	0.32 – 2.96	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.97	0.41 – 5.75	4
Kidney & Renal Pelvis	2.4	17.2	14.1	14.1	13.6	1.21	0.63 – 2.12	4
Larynx	<1	*	4.7	4.4	3.5	0.78	0.09 – 2.82	3
Leukemia	2.2	13.4	11.8	11.1	12.2	1.25	0.62 – 2.23	3
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.62	0.13 – 1.81	3
Lung & Bronchus	11.4	74.2	70.7	75.0	63.1	1.15	0.87 – 1.49	4
Melanoma of the Skin	3.2	19.1	15.3	17.6	19.6	1.19	0.68 – 1.93	2
Multiple Myeloma	1.4	9.7	6.0	5.1	5.6	1.60	0.64 – 3.30	4
Non-Hodgkin's Lymphoma	3.4	23.0	19.1	19.2	19.5	1.18	0.69 – 1.89	3
Oral Cavity & Pharynx	1.4	10.8	10.4	9.6	10.4	0.95	0.38 – 1.96	3
Ovary	<1	*	12.3	12.3	13.1	0.76	0.21 – 1.94	2
Pancreas	2.0	12.7	12.3	10.9	11.7	1.08	0.52 – 1.99	4
Prostate (Male)	8.0	125.5	157.5	144.7	159.3	0.79	0.56 – 1.07	1
Stomach	1.2	8.1	8.1	6.1	7.9	0.98	0.36 – 2.12	4
Testis (Male)	<1	*	4.9	5.5	5.4	0.8	0.02 – 4.48	1
Thyroid	1.4	12.8	7.9	8.3	9.6	1.23	0.49 – 2.53	4

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Orange

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	17.0	432.3	477.1	466.4	462.9	0.90	0.72 – 1.11	1
Bladder	0.0	*	21.5	21.6	21.0	0.23	0.01 – 1.29	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.83	0.02 – 4.63	3
Breast (Female)	2.0	97.5	122.2	120.4	123.8	0.76	0.36 – 1.39	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.8	46.9	51.1	52.1	49.1	0.9	0.41 – 1.70	4
Corpus Uterus (Female)	1.2	47.4	29.9	26.4	23.3	2.43	0.89 – 5.29	4
Esophagus	<1	*	5.7	5.5	4.5	1.07	0.03 – 5.98	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.38	0.06 – 13.24	4
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.72	0.09 – 2.61	2
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	0.42	0.01 – 2.36	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	3.0	76.2	70.7	75.0	63.1	1.14	0.64 – 1.88	4
Melanoma of the Skin	1.2	26.8	15.3	17.6	19.6	1.58	0.58 – 3.44	2
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.72	0.21 – 6.20	4
Non-Hodgkin's Lymphoma	1.2	27.4	19.1	19.2	19.5	1.54	0.56 – 3.34	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.95	0.12 – 3.44	4
Ovary	<1	*	12.3	12.3	13.1	0.73	0.02 – 4.04	2
Pancreas	<1	*	12.3	10.9	11.7	0.84	0.10 – 3.02	3
Prostate (Male)	2.4	130.0	157.5	144.7	159.3	0.77	0.40 – 1.35	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	0.58	0.01 – 3.25	2

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Parma

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	476.0	421.6	477.1	466.4	462.9	0.90	0.86 – 0.94	1
Bladder	29.0	24.0	21.5	21.6	21.0	1.09	0.92 – 1.28	3
Brain and Other CNS	4.6	4.7	5.8	6.8	6.4	0.70	0.45 – 1.05	2
Breast (Female)	59.4	100.5	122.2	120.4	123.8	0.81	0.72 – 0.91	1
Cervix (Female)	3.4	7.2	8.0	8.0	8.2	0.83	0.49 – 1.33	4
Colon & Rectum	53.6	44.2	51.1	52.1	49.1	0.91	0.80 – 1.02	1
Corpus Uterus (Female)	18.8	31.5	29.9	26.4	23.3	1.35	1.09 – 1.65	3
Esophagus	4.8	4.1	5.7	5.5	4.5	0.90	0.58 – 1.34	2
Hodgkin's Lymphoma	1.6	2.1	2.9	3.0	2.8	0.64	0.28 – 1.26	2
Kidney & Renal Pelvis	13.0	11.9	14.1	14.1	13.6	0.86	0.66 – 1.09	3
Larynx	4.4	3.9	4.7	4.4	3.5	1.13	0.71 – 1.72	4
Leukemia	14.4	13.0	11.8	11.1	12.2	1.04	0.82 – 1.31	3
Liver & Intrahepatic Bile Duct	3.6	3.2	5.6	4.3	6.6	0.49	0.29 – 0.77	3
Lung & Bronchus	69.4	57.6	70.7	75.0	63.1	0.90	0.81 – 1.00	3
Melanoma of the Skin	19.4	18.9	15.3	17.6	19.6	0.93	0.75 – 1.14	2
Multiple Myeloma	6.2	5.1	6.0	5.1	5.6	0.92	0.62 – 1.30	3
Non-Hodgkin's Lymphoma	21.2	18.4	19.1	19.2	19.5	0.95	0.78 – 1.15	2
Oral Cavity & Pharynx	9.2	8.6	10.4	9.6	10.4	0.83	0.60 – 1.10	3
Ovary	9.0	16.6	12.3	12.3	13.1	1.12	0.81 – 1.50	4
Pancreas	14.6	12.1	12.3	10.9	11.7	1.02	0.80 – 1.28	3
Prostate (Male)	59.8	124.7	157.5	144.7	159.3	0.77	0.68 – 0.86	1
Stomach	6.6	5.6	8.1	6.1	7.9	0.69	0.48 – 0.97	2
Testis (Male)	2.6	6.3	4.9	5.5	5.4	1.19	0.63 – 2.03	2
Thyroid	7.4	7.5	7.9	8.3	9.6	0.83	0.58 – 1.14	2

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Parma Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	157.0	475.0	477.1	466.4	462.9	0.99	0.92 – 1.06	3
Bladder	11.0	32.2	21.5	21.6	21.0	1.35	1.02 – 1.75	4
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.43	0.12 – 1.11	2
Breast (Female)	21.4	129.0	122.2	120.4	123.8	0.97	0.80 – 1.17	3
Cervix (Female)	1.8	15.5	8.0	8.0	8.2	1.62	0.74 – 3.08	4
Colon & Rectum	14.4	39.0	51.1	52.1	49.1	0.78	0.61 – 0.98	1
Corpus Uterus (Female)	6.4	38.1	29.9	26.4	23.3	1.53	1.05 – 2.16	4
Esophagus	1.8	5.5	5.7	5.5	4.5	1.11	0.51 – 2.11	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	1.21	0.33 – 3.09	4
Kidney & Renal Pelvis	3.0	9.3	14.1	14.1	13.6	0.67	0.37 – 1.10	1
Larynx	2.6	6.9	4.7	4.4	3.5	2.27	1.21 – 3.88	4
Leukemia	3.8	14.9	11.8	11.1	12.2	0.91	0.55 – 1.41	4
Liver & Intrahepatic Bile Duct	1.0	1.9	5.6	4.3	6.6	0.46	0.15 – 1.07	1
Lung & Bronchus	27.0	71.4	70.7	75.0	63.1	1.13	0.95 – 1.34	3
Melanoma of the Skin	3.4	10.8	15.3	17.6	19.6	0.56	0.33 – 0.90	2
Multiple Myeloma	2.8	7.4	6.0	5.1	5.6	1.34	0.73 – 2.25	4
Non-Hodgkin's Lymphoma	7.4	20.6	19.1	19.2	19.5	1.09	0.77 – 1.51	4
Oral Cavity & Pharynx	4.4	14.9	10.4	9.6	10.4	1.36	0.85 – 2.06	4
Ovary	2.0	13.3	12.3	12.3	13.1	0.81	0.39 – 1.48	3
Pancreas	3.4	8.3	12.3	10.9	11.7	0.76	0.44 – 1.21	1
Prostate (Male)	17.8	128.9	157.5	144.7	159.3	0.81	0.65 – 1.00	2
Stomach	3.2	8.5	8.1	6.1	7.9	1.08	0.62 – 1.75	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.75	0.09 – 2.72	3
Thyroid	1.8	7.9	7.9	8.3	9.6	0.76	0.35 – 1.45	4

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Pepper Pike

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	36.0	413.1	477.1	466.4	462.9	0.85	0.73 – 0.99	1
Bladder	2.0	26.1	21.5	21.6	21.0	1.11	0.56 – 1.99	3
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	5.2	108.5	122.2	120.4	123.8	0.89	0.58 – 1.30	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	3.0	35.1	51.1	52.1	49.1	0.66	0.37 – 1.09	2
Corpus Uterus (Female)	1.4	27.5	29.9	26.4	23.3	1.23	0.49 – 2.53	2
Esophagus	<1	*	5.7	5.5	4.5	0.46	0.01 – 2.58	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.42	0.29 – 8.73	4
Kidney & Renal Pelvis	1.2	12.5	14.1	14.1	13.6	0.96	0.35 – 2.08	3
Larynx	<1	*	4.7	4.4	3.5	1.19	0.14 – 4.29	3
Leukemia	1.2	15.2	11.8	11.1	12.2	1.17	0.43 – 2.55	3
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.32	0.01 – 1.78	2
Lung & Bronchus	3.0	33.4	70.7	75.0	63.1	0.49	0.28 – 0.81	1
Melanoma of the Skin	1.8	23.1	15.3	17.6	19.6	1.10	0.50 – 2.08	1
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.38	0.01 – 2.10	2
Non-Hodgkin's Lymphoma	2.0	23.1	19.1	19.2	19.5	1.15	0.55 – 2.12	3
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.42	0.05 – 1.52	2
Ovary	<1	*	12.3	12.3	13.1	1.30	0.35 – 3.32	4
Pancreas	<1	*	12.3	10.9	11.7	0.73	0.20 – 1.87	2
Prostate (Male)	8.4	177.1	157.5	144.7	159.3	1.11	0.80 – 1.51	4
Stomach	<1	*	8.1	6.1	7.9	0.28	0.01 – 1.54	2
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	<1	*	7.9	8.3	9.6	1.16	0.32 – 2.97	4

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Richmond Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	63.0	451.1	477.1	466.4	462.9	0.96	0.86 – 1.08	3
Bladder	4.0	24.7	21.5	21.6	21.0	1.21	0.73 – 1.89	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	0.73	0.15 – 2.14	2
Breast (Female)	9.4	119.7	122.2	120.4	123.8	1.00	0.74 – 1.33	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.36	0.01 – 2.03	3
Colon & Rectum	6.8	48	51.1	52.1	49.1	0.94	0.65 – 1.32	2
Corpus Uterus (Female)	3.2	43.9	29.9	26.4	23.3	1.81	1.03 – 2.94	4
Esophagus	<1	*	5.7	5.5	4.5	0.61	0.07 – 2.22	1
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.62	0.02 – 3.46	2
Kidney & Renal Pelvis	3.4	24.2	14.1	14.1	13.6	1.80	1.05 – 2.88	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	1.2	9	11.8	11.1	12.2	0.71	0.26 – 1.56	1
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.43	0.05 – 1.56	2
Lung & Bronchus	6.8	48.5	70.7	75.0	63.1	0.73	0.51 – 1.02	2
Melanoma of the Skin	1.8	13.3	15.3	17.6	19.6	0.68	0.31 – 1.29	1
Multiple Myeloma	1.0	6.8	6.0	5.1	5.6	1.21	0.39 – 2.83	4
Non-Hodgkin's Lymphoma	3.0	20.1	19.1	19.2	19.5	1.09	0.61 – 1.80	2
Oral Cavity & Pharynx	1.4	11.2	10.4	9.6	10.4	0.99	0.40 – 2.03	4
Ovary	<1	*	12.3	12.3	13.1	0.79	0.21 – 2.02	2
Pancreas	1.6	11.0	12.3	10.9	11.7	0.93	0.40 – 1.82	3
Prostate (Male)	9.4	158.7	157.5	144.7	159.3	0.96	0.71 – 1.28	3
Stomach	1.6	10.9	8.1	6.1	7.9	1.38	0.60 – 2.72	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.43	0.17 – 5.16	3
Thyroid	<1	*	7.9	8.3	9.6	0.68	0.19 – 1.74	2

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Rocky River

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	152.0	456.4	477.1	466.4	462.9	0.99	0.93 – 1.07	3
Bladder	8.0	21.2	21.5	21.6	21.0	1.07	0.77 – 1.45	2
Brain and Other CNS	1.6	6.4	5.8	6.8	6.4	0.90	0.39 – 1.77	3
Breast (Female)	22.4	129.4	122.2	120.4	123.8	1.03	0.85 – 1.24	3
Cervix (Female)	1.0	8.4	8.0	8.0	8.2	0.90	0.29 – 2.10	4
Colon & Rectum	18.0	49	51.1	52.1	49.1	1.02	0.82 – 1.26	3
Corpus Uterus (Female)	4.6	29.2	29.9	26.4	23.3	1.11	0.70 – 1.67	2
Esophagus	2.6	7.2	5.7	5.5	4.5	1.67	0.89 – 2.85	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.66	0.08 – 2.38	3
Kidney & Renal Pelvis	3.4	9.6	14.1	14.1	13.6	0.78	0.46 – 1.25	2
Larynx	1.0	3.2	4.7	4.4	3.5	0.89	0.29 – 2.08	4
Leukemia	6.4	17.8	11.8	11.1	12.2	1.60	1.10 – 2.26	4
Liver & Intrahepatic Bile Duct	1.0	2.8	5.6	4.3	6.6	0.47	0.15 – 1.09	3
Lung & Bronchus	17.6	48.9	70.7	75.0	63.1	0.78	0.63 – 0.96	2
Melanoma of the Skin	9.4	29.9	15.3	17.6	19.6	1.61	1.18 – 2.14	4
Multiple Myeloma	1.6	4.9	6.0	5.1	5.6	0.80	0.35 – 1.58	2
Non-Hodgkin's Lymphoma	8.6	25	19.1	19.2	19.5	1.33	0.96 – 1.79	3
Oral Cavity & Pharynx	2.4	8.6	10.4	9.6	10.4	0.75	0.39 – 1.31	2
Ovary	1.8	11.3	12.3	12.3	13.1	0.74	0.34 – 1.41	2
Pancreas	3.2	9.7	12.3	10.9	11.7	0.75	0.43 – 1.21	2
Prostate (Male)	18.4	137.7	157.5	144.7	159.3	0.87	0.70 – 1.06	2
Stomach	2.6	6.3	8.1	6.1	7.9	0.92	0.49 – 1.57	3
Testis (Male)	<1	*	4.9	5.5	5.4	0.89	0.11 – 3.22	2
Thyroid	1.6	6.7	7.9	8.3	9.6	0.70	0.30 – 1.37	2

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Seven Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	98.0	484.8	477.1	466.4	462.9	1.02	0.93 – 1.12	4
Bladder	6.0	27.2	21.5	21.6	21.0	1.22	0.82 – 1.76	4
Brain and Other CNS	1.4	9.3	5.8	6.8	6.4	1.27	0.51 – 2.62	4
Breast (Female)	11.6	111.4	122.2	120.4	123.8	0.93	0.70 – 1.20	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.63	0.08 – 2.28	4
Colon & Rectum	9.4	46.2	51.1	52.1	49.1	0.89	0.65 – 1.18	2
Corpus Uterus (Female)	4.8	40.5	29.9	26.4	23.3	1.96	1.25 – 2.91	4
Esophagus	1.6	8.0	5.7	5.5	4.5	1.62	0.70 – 3.20	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	2.21	0.6 – 5.65	4
Kidney & Renal Pelvis	1.4	7.1	14.1	14.1	13.6	0.50	0.20 – 1.04	1
Larynx	1.8	8.7	4.7	4.4	3.5	2.44	1.12 – 4.64	4
Leukemia	3.0	17.7	11.8	11.1	12.2	1.27	0.71 – 2.10	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.44	0.09 – 1.29	2
Lung & Bronchus	12.2	52.6	70.7	75.0	63.1	0.85	0.65 – 1.09	3
Melanoma of the Skin	2.0	13.6	15.3	17.6	19.6	0.55	0.27 – 1.02	2
Multiple Myeloma	1.2	4.7	6.0	5.1	5.6	0.97	0.36 – 2.12	2
Non-Hodgkin's Lymphoma	5.6	25.5	19.1	19.2	19.5	1.41	0.94 – 2.04	4
Oral Cavity & Pharynx	2.0	9.2	10.4	9.6	10.4	0.99	0.47 – 1.82	3
Ovary	1.4	14.8	12.3	12.3	13.1	1.03	0.41 – 2.12	4
Pancreas	1.6	7.2	12.3	10.9	11.7	0.62	0.27 – 1.22	1
Prostate (Male)	16.6	164.9	157.5	144.7	159.3	1.01	0.80 – 1.25	4
Stomach	2.2	9.9	8.1	6.1	7.9	1.30	0.65 – 2.32	4
Testis (Male)	<1	*	4.9	5.5	5.4	3.01	0.82 – 7.69	4
Thyroid	1.6	10.4	7.9	8.3	9.6	1.12	0.48 – 2.21	3

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Shaker Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	151.0	437.1	477.1	466.4	462.9	0.94	0.87 – 1.01	1
Bladder	6.0	16.4	21.5	21.6	21.0	0.80	0.54 – 1.14	2
Brain and Other CNS	1.8	6.1	5.8	6.8	6.4	0.86	0.39 – 1.63	3
Breast (Female)	30.4	155.5	122.2	120.4	123.8	1.26	1.06 – 1.47	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.41	0.09 – 1.21	2
Colon & Rectum	14.6	42.1	51.1	52.1	49.1	0.84	0.66 – 1.05	2
Corpus Uterus (Female)	6.0	28.9	29.9	26.4	23.3	1.31	0.88 – 1.87	2
Esophagus	1.2	3.5	5.7	5.5	4.5	0.75	0.28 – 1.64	1
Hodgkin's Lymphoma	1.2	3.9	2.9	3.0	2.8	1.50	0.55 – 3.26	3
Kidney & Renal Pelvis	3.6	10.2	14.1	14.1	13.6	0.77	0.45 – 1.21	1
Larynx	<1	*	4.7	4.4	3.5	0.66	0.18 – 1.69	3
Leukemia	4.8	13.5	11.8	11.1	12.2	1.15	0.74 – 1.72	3
Liver & Intrahepatic Bile Duct	1.2	3.4	5.6	4.3	6.6	0.52	0.19 – 1.13	3
Lung & Bronchus	13.6	38.0	70.7	75.0	63.1	0.61	0.47 – 0.77	1
Melanoma of the Skin	6.4	19.7	15.3	17.6	19.6	0.97	0.67 – 1.38	3
Multiple Myeloma	2.4	7.3	6.0	5.1	5.6	1.20	0.62 – 2.10	3
Non-Hodgkin's Lymphoma	6.0	17.4	19.1	19.2	19.5	0.89	0.60 – 1.27	3
Oral Cavity & Pharynx	1.6	4.8	10.4	9.6	10.4	0.45	0.19 – 0.88	1
Ovary	1.8	9.1	12.3	12.3	13.1	0.70	0.32 – 1.32	2
Pancreas	5.2	14.5	12.3	10.9	11.7	1.25	0.82 – 1.83	4
Prostate (Male)	24.8	169.9	157.5	144.7	159.3	1.06	0.88 – 1.27	2
Stomach	1.8	5.1	8.1	6.1	7.9	0.64	0.29 – 1.22	2
Testis (Male)	1.2	11.3	4.9	5.5	5.4	1.85	0.68 – 4.02	4
Thyroid	2.4	7.4	7.9	8.3	9.6	0.80	0.41 – 1.39	3

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Solon

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	112.0	562.3	477.1	466.4	462.9	1.16	1.07 – 1.26	4
Bladder	6.0	34.2	21.5	21.6	21.0	1.55	1.06 – 2.19	4
Brain and Other CNS	2.6	11.7	5.8	6.8	6.4	1.88	1.00 – 3.21	4
Breast (Female)	21.8	187.7	122.2	120.4	123.8	1.49	1.22 – 1.79	4
Cervix (Female)	<1	*	8.0	8.0	8.2	0.41	0.05 – 1.48	4
Colon & Rectum	10.4	57.9	51.1	52.1	49.1	1.05	0.78 – 1.37	4
Corpus Uterus (Female)	3.0	25.6	29.9	26.4	23.3	1.12	0.63 – 1.84	2
Esophagus	<1	*	5.7	5.5	4.5	0.86	0.23 – 2.20	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.36	0.01 – 2.01	3
Kidney & Renal Pelvis	3.4	17.5	14.1	14.1	13.6	1.16	0.68 – 1.86	4
Larynx	<1	*	4.7	4.4	3.5	0.80	0.17 – 2.34	3
Leukemia	3.0	16.7	11.8	11.1	12.2	1.21	0.68 – 2.00	4
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.28	0.03 – 1.00	2
Lung & Bronchus	11.0	55.4	70.7	75.0	63.1	0.86	0.65 – 1.12	3
Melanoma of the Skin	5.4	27.4	15.3	17.6	19.6	1.27	0.84 – 1.85	2
Multiple Myeloma	1.4	6.8	6.0	5.1	5.6	1.21	0.49 – 2.50	4
Non-Hodgkin's Lymphoma	4.4	22.8	19.1	19.2	19.5	1.08	0.68 – 1.64	2
Oral Cavity & Pharynx	2.0	9.4	10.4	9.6	10.4	0.87	0.42 – 1.60	4
Ovary	1.6	14.6	12.3	12.3	13.1	1.07	0.46 – 2.11	4
Pancreas	2.4	11.9	12.3	10.9	11.7	1.04	0.54 – 1.81	2
Prostate (Male)	18.0	183.7	157.5	144.7	159.3	1.18	0.95 – 1.45	4
Stomach	2.0	10.9	8.1	6.1	7.9	1.25	0.60 – 2.30	4
Testis (Male)	<1	*	4.9	5.5	5.4	1.63	0.44 – 4.16	3
Thyroid	2.2	11.6	7.9	8.3	9.6	1.02	0.51 – 1.82	3

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: South Euclid

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	114.0	434.3	477.1	466.4	462.9	0.92	0.85 – 1.00	2
Bladder	5.0	17.3	21.5	21.6	21.0	0.81	0.52 – 1.21	1
Brain and Other CNS	1.0	4.1	5.8	6.8	6.4	0.61	0.20 – 1.43	1
Breast (Female)	20.2	136.2	122.2	120.4	123.8	1.09	0.89 – 1.32	4
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	14.0	48.7	51.1	52.1	49.1	1.03	0.80 – 1.30	3
Corpus Uterus (Female)	4.4	30.6	29.9	26.4	23.3	1.28	0.80 – 1.93	2
Esophagus	1.0	4.0	5.7	5.5	4.5	0.82	0.27 – 1.92	2
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.61	0.07 – 2.19	1
Kidney & Renal Pelvis	2.0	7.6	14.1	14.1	13.6	0.56	0.27 – 1.03	1
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	4.4	17.5	11.8	11.1	12.2	1.34	0.84 – 2.03	4
Liver & Intrahepatic Bile Duct	1.4	5.9	5.6	4.3	6.6	0.8	0.32 – 1.65	4
Lung & Bronchus	14.2	52.5	70.7	75.0	63.1	0.82	0.64 – 1.03	3
Melanoma of the Skin	3.0	11.3	15.3	17.6	19.6	0.59	0.33 – 0.97	1
Multiple Myeloma	1.8	6.9	6.0	5.1	5.6	1.16	0.53 – 2.21	4
Non-Hodgkin's Lymphoma	4.8	18.1	19.1	19.2	19.5	0.91	0.58 – 1.36	1
Oral Cavity & Pharynx	2.4	8.6	10.4	9.6	10.4	0.89	0.46 – 1.56	2
Ovary	<1	*	12.3	12.3	13.1	0.40	0.11 – 1.03	1
Pancreas	4.0	14.3	12.3	10.9	11.7	1.23	0.75 – 1.90	4
Prostate (Male)	15.6	151.6	157.5	144.7	159.3	0.91	0.72 – 1.13	3
Stomach	2.0	7.9	8.1	6.1	7.9	0.91	0.44 – 1.68	3
Testis (Male)	<1	*	4.9	5.5	5.4	1.04	0.22 – 3.05	2
Thyroid	2.2	9.3	7.9	8.3	9.6	0.93	0.47 – 1.67	3

Peer Comparison Neighborhoods/Municipalities: Bay Village, Broadview Heights, Brooklyn Heights, Chagrin Falls Village, Fairview Park, Mayfield Village, Middleburg Heights, Olmsted Township, Parma, Richmond Heights, Rocky River, Seven Hills, South Euclid

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Strongsville

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	218.0	502.0	477.1	466.4	462.9	1.06	1.00 – 1.13	3
Bladder	13.0	31.2	21.5	21.6	21.0	1.45	1.12 – 1.86	4
Brain and Other CNS	2.4	5.4	5.8	6.8	6.4	0.84	0.43 – 1.46	2
Breast (Female)	31.2	124.8	122.2	120.4	123.8	1.01	0.86 – 1.18	2
Cervix (Female)	1.0	3.9	8.0	8.0	8.2	0.50	0.16 – 1.16	2
Colon & Rectum	21.4	51.8	51.1	52.1	49.1	1.01	0.83 – 1.23	3
Corpus Uterus (Female)	7.4	30.3	29.9	26.4	23.3	1.28	0.90 – 1.76	3
Esophagus	2.4	5.7	5.7	5.5	4.5	1.21	0.62 – 2.11	3
Hodgkin's Lymphoma	1.0	2.6	2.9	3.0	2.8	0.86	0.28 – 2	3
Kidney & Renal Pelvis	7.4	16.6	14.1	14.1	13.6	1.20	0.85 – 1.66	4
Larynx	2.6	5.1	4.7	4.4	3.5	1.64	0.87 – 2.80	4
Leukemia	6.0	14.5	11.8	11.1	12.2	1.16	0.78 – 1.65	3
Liver & Intrahepatic Bile Duct	2.2	5.4	5.6	4.3	6.6	0.72	0.36 – 1.30	3
Lung & Bronchus	23.0	53.7	70.7	75.0	63.1	0.85	0.70 – 1.01	3
Melanoma of the Skin	8.6	19.3	15.3	17.6	19.6	0.97	0.70 – 1.31	2
Multiple Myeloma	2.4	5.7	6.0	5.1	5.6	0.98	0.51 – 1.71	4
Non-Hodgkin's Lymphoma	11.6	26.6	19.1	19.2	19.5	1.36	1.03 – 1.76	4
Oral Cavity & Pharynx	3.4	8.0	10.4	9.6	10.4	0.71	0.41 – 1.13	2
Ovary	4.0	15.6	12.3	12.3	13.1	1.25	0.76 – 1.93	2
Pancreas	6.0	14.2	12.3	10.9	11.7	1.21	0.82 – 1.73	2
Prostate (Male)	35.4	192.3	157.5	144.7	159.3	1.12	0.96 – 1.29	4
Stomach	3.4	9.2	8.1	6.1	7.9	1.00	0.58 – 1.61	4
Testis (Male)	1.0	4.5	4.9	5.5	5.4	0.91	0.30 – 2.13	2
Thyroid	4.8	10	7.9	8.3	9.6	1.07	0.69 – 1.59	2

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: University Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	61.0	449.5	477.1	466.4	462.9	0.94	0.84 – 1.05	2
Bladder	4.0	26.3	21.5	21.6	21.0	1.29	0.79 – 2.00	3
Brain and Other CNS	1.0	7.8	5.8	6.8	6.4	1.12	0.36 – 2.60	4
Breast (Female)	9.0	116.7	122.2	120.4	123.8	0.97	0.71 – 1.30	2
Cervix (Female)	<1	*	8.0	8.0	8.2	0.68	0.08 – 2.47	3
Colon & Rectum	6.4	48.8	51.1	52.1	49.1	0.90	0.62 – 1.28	3
Corpus Uterus (Female)	2.4	31.7	29.9	26.4	23.3	1.38	0.71 – 2.41	3
Esophagus	<1	*	5.7	5.5	4.5	0.63	0.08 – 2.27	2
Hodgkin's Lymphoma	1	6.1	2.9	3.0	2.8	2.41	0.78 – 5.63	4
Kidney & Renal Pelvis	1.4	10.3	14.1	14.1	13.6	0.75	0.30 – 1.54	2
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	1.4	11.3	11.8	11.1	12.2	0.79	0.32 – 1.63	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	0.43	0.05 – 1.57	3
Lung & Bronchus	7.0	51.3	70.7	75.0	63.1	0.77	0.53 – 1.07	1
Melanoma of the Skin	2.8	20.5	15.3	17.6	19.6	1.03	0.56 – 1.73	3
Multiple Myeloma	<1	*	6.0	5.1	5.6	0.99	0.27 – 2.53	2
Non-Hodgkin's Lymphoma	1.8	13.2	19.1	19.2	19.5	0.65	0.30 – 1.23	2
Oral Cavity & Pharynx	1.4	10.8	10.4	9.6	10.4	0.99	0.40 – 2.05	4
Ovary	1.0	10.1	12.3	12.3	13.1	0.98	0.32 – 2.29	2
Pancreas	1.4	10.8	12.3	10.9	11.7	0.83	0.33 – 1.70	2
Prostate (Male)	9.6	173.8	157.5	144.7	159.3	1.02	0.75 – 1.35	3
Stomach	<1	*	8.1	6.1	7.9	0.53	0.11 – 1.54	1
Testis (Male)	<1	*	4.9	5.5	5.4	0.53	0.01 – 2.95	3
Thyroid	2.0	16.1	7.9	8.3	9.6	1.55	0.74 – 2.85	4

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Valley View

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	13.0	537.8	477.1	466.4	462.9	1.15	0.89 – 1.47	4
Bladder	1.0	29.7	21.5	21.6	21.0	1.21	0.25 – 3.52	3
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.32	0.03 – 7.33	4
Breast (Female)	1.8	134.5	122.2	120.4	123.8	1.07	0.49 – 2.04	2
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	1.0	36.4	51.1	52.1	49.1	0.84	0.27 – 1.97	2
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.90	0.39 – 5.56	4
Esophagus	<1	*	5.7	5.5	4.5	3.55	0.43 – 12.8	4
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	0.58	0.01 – 3.24	3
Larynx	<1	*	4.7	4.4	3.5	4.41	0.53 – 15.9	4
Leukemia	<1	*	11.8	11.1	12.2	1.43	0.17 – 5.16	2
Liver & Intrahepatic Bile Duct	<1	*	5.6	4.3	6.6	1.17	0.03 – 6.54	4
Lung & Bronchus	1.8	80.7	70.7	75.0	63.1	1.16	0.53 – 2.20	4
Melanoma of the Skin	1.0	35.5	15.3	17.6	19.6	2.08	0.67 – 4.85	3
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	<1	*	19.1	19.2	19.5	1.27	0.26 – 3.71	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.75	0.02 – 4.16	2
Ovary	<1	*	12.3	12.3	13.1	1.16	0.03 – 6.44	3
Pancreas	<1	*	12.3	10.9	11.7	1.43	0.17 – 5.18	4
Prostate (Male)	2	159	157.5	144.7	159.3	1.05	0.51 – 1.94	3
Stomach	<1	*	8.1	6.1	7.9	2.11	0.26 – 7.61	4
Testis (Male)	<1	*	4.9	5.5	5.4	3.83	0.1 – 21.36	4
Thyroid	<1	*	7.9	8.3	9.6	0.85	0.02 – 4.76	2

Peer Comparison Neighborhoods/Municipalities: Beachwood, Bratenahl, Brecksville, Chagrin Falls Township, Hunting Valley, Moreland Hills, Orange, Pepper Pike, Solon, Valley View

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Walton Hills

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	22.0	484.9	477.1	466.4	462.9	1.10	0.90 – 1.33	4
Bladder	1.0	11.9	21.5	21.6	21.0	0.60	0.12 – 1.75	1
Brain and Other CNS	<1	*	5.8	6.8	6.4	1.78	0.22 – 6.44	3
Breast (Female)	1.4	64.6	122.2	120.4	123.8	0.53	0.21 – 1.09	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	3.8	81.9	51.1	52.1	49.1	1.69	1.02 – 2.64	4
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	1.18	0.24 – 3.45	3
Esophagus	<1	*	5.7	5.5	4.5	0.98	0.02 – 5.46	3
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	1.2	28.1	14.1	14.1	13.6	2.10	0.77 – 4.58	4
Larynx	0.0	*	4.7	4.4	3.5	0.00	0.00 – 0.00	1
Leukemia	<1	*	11.8	11.1	12.2	1.20	0.25 – 3.52	4
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	3.2	70.8	70.7	75.0	63.1	1.08	0.62 – 1.76	3
Melanoma of the Skin	1.0	23.4	15.3	17.6	19.6	1.34	0.43 – 3.12	4
Multiple Myeloma	<1	*	6.0	5.1	5.6	1.55	0.19 – 5.61	4
Non-Hodgkin's Lymphoma	1.4	27.7	19.1	19.2	19.5	1.69	0.68 – 3.48	4
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	1.43	0.30 – 4.19	4
Ovary	<1	*	12.3	12.3	13.1	1.37	0.17 – 4.96	3
Pancreas	<1	*	12.3	10.9	11.7	0.73	0.09 – 2.64	3
Prostate (Male)	3.8	187.7	157.5	144.7	159.3	1.17	0.71 – 1.83	3
Stomach	<1	*	8.1	6.1	7.9	1.67	0.34 – 4.87	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Bedford, Bedford Heights, Berea, Brooklyn, Cuyahoga Heights, Kamms Corners, Maple Heights, Mayfield Heights, Oakwood, Parma Heights, Shaker Heights, University Heights, Walton Hills

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Warrensville Heights

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	79.0	454.1	477.1	466.4	462.9	1.00	0.90 – 1.10	1
Bladder	3.0	18.9	21.5	21.6	21.0	0.88	0.50 – 1.43	1
Brain and Other CNS	1.2	6.8	5.8	6.8	6.4	1.15	0.42 – 2.49	3
Breast (Female)	9.2	91.6	122.2	120.4	123.8	0.73	0.53 – 0.97	1
Cervix (Female)	<1	*	8.0	8.0	8.2	0.26	0.01 – 1.44	1
Colon & Rectum	10.6	59.5	51.1	52.1	49.1	1.25	0.94 – 1.64	4
Corpus Uterus (Female)	1.2	11.0	29.9	26.4	23.3	0.49	0.18 – 1.08	1
Esophagus	1.2	6.6	5.7	5.5	4.5	1.53	0.56 – 3.32	3
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.48	0.01 – 2.67	1
Kidney & Renal Pelvis	2.6	15.0	14.1	14.1	13.6	1.12	0.59 – 1.91	2
Larynx	<1	*	4.7	4.4	3.5	1.31	0.36 – 3.37	2
Leukemia	1.8	10.8	11.8	11.1	12.2	0.88	0.40 – 1.68	3
Liver & Intrahepatic Bile Duct	1.2	7.1	5.6	4.3	6.6	1.05	0.38 – 2.28	4
Lung & Bronchus	11.4	65.9	70.7	75.0	63.1	1.04	0.79 – 1.35	1
Melanoma of the Skin	<1	*	15.3	17.6	19.6	0.12	0.01 – 0.45	1
Multiple Myeloma	1.6	9.2	6.0	5.1	5.6	1.64	0.71 – 3.24	4
Non-Hodgkin's Lymphoma	2.0	12.2	19.1	19.2	19.5	0.61	0.29 – 1.11	1
Oral Cavity & Pharynx	<1	*	10.4	9.6	10.4	0.22	0.03 – 0.81	1
Ovary	1.0	8.8	12.3	12.3	13.1	0.74	0.24 – 1.73	1
Pancreas	2.2	12.6	12.3	10.9	11.7	1.09	0.54 – 1.95	2
Prostate (Male)	17.6	239.3	157.5	144.7	159.3	1.56	1.25 – 1.92	4
Stomach	1.6	9.7	8.1	6.1	7.9	1.18	0.51 – 2.32	3
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	1.6	10.0	7.9	8.3	9.6	1.06	0.46 – 2.09	4

Peer Comparison Neighborhoods/Municipalities: Brookpark, Cleveland Heights, Garfield Heights, Glenwillow, Jefferson, Lakewood, Lee-Miles, Newburgh Heights, Old Brooklyn, Warrensville Heights

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Westlake

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	193.0	464.6	477.1	466.4	462.9	0.99	0.93 – 1.06	2
Bladder	11.0	23.3	21.5	21.6	21.0	1.11	0.83 – 1.45	3
Brain and Other CNS	2.2	6.0	5.8	6.8	6.4	0.91	0.46 – 1.63	2
Breast (Female)	30.2	134	122.2	120.4	123.8	1.08	0.92 – 1.27	3
Cervix (Female)	2.0	11.0	8.0	8.0	8.2	1.28	0.61 – 2.35	4
Colon & Rectum	20.4	43.6	51.1	52.1	49.1	0.93	0.75 – 1.12	1
Corpus Uterus (Female)	3.8	15.5	29.9	26.4	23.3	0.73	0.44 – 1.14	1
Esophagus	3.2	7.1	5.7	5.5	4.5	1.65	0.94 – 2.68	4
Hodgkin's Lymphoma	<1	*	2.9	3.0	2.8	0.89	0.24 – 2.28	4
Kidney & Renal Pelvis	4.8	11.9	14.1	14.1	13.6	0.86	0.55 – 1.28	1
Larynx	1.2	3.1	4.7	4.4	3.5	0.85	0.31 – 1.84	3
Leukemia	5.0	11.3	11.8	11.1	12.2	0.98	0.63 – 1.44	2
Liver & Intrahepatic Bile Duct	2.2	5.3	5.6	4.3	6.6	0.80	0.40 – 1.43	2
Lung & Bronchus	20.6	48.6	70.7	75.0	63.1	0.75	0.62 – 0.91	2
Melanoma of the Skin	13.4	36.9	15.3	17.6	19.6	1.72	1.33 – 2.18	4
Multiple Myeloma	1.6	3.9	6.0	5.1	5.6	0.65	0.28 – 1.27	2
Non-Hodgkin's Lymphoma	6.6	15.5	19.1	19.2	19.5	0.80	0.55 – 1.12	1
Oral Cavity & Pharynx	4.6	11.0	10.4	9.6	10.4	1.09	0.69 – 1.63	4
Ovary	4.0	18.0	12.3	12.3	13.1	1.30	0.80 – 2.01	3
Pancreas	5.4	12.4	12.3	10.9	11.7	1.02	0.67 – 1.48	1
Prostate (Male)	24.2	140.2	157.5	144.7	159.3	0.88	0.73 – 1.05	1
Stomach	2.0	4.7	8.1	6.1	7.9	0.56	0.27 – 1.04	1
Testis (Male)	1.0	7.9	4.9	5.5	5.4	1.34	0.43 – 3.12	4
Thyroid	4.0	11.3	7.9	8.3	9.6	1.19	0.73 – 1.83	3

Peer Comparison Neighborhoods/Municipalities: Bentleyville, Gates Mills, Highland Heights, Independence, Lyndhurst, North Olmsted, North Royalton, Olmsted Falls, Strongsville, Westlake

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.

City at a Glance: Woodmere

Average Annual Age-Adjusted Incidence Rates from 2002-2006 (per 100,000 people)

Primary Cancer Site/Type	Average Annual Cases 2002-2006	City Rate	Cuyahoga Rate	Ohio Rate	National Rate	Standardized Incidence Ratio (SIR) ¹	SIR 95% Confidence Interval	Peer Comparison Rank
All Sites/Types	3.0	345.6	477.1	466.4	462.9	0.72	0.39 – 1.24	1
Bladder	0.0	*	21.5	21.6	21.0	1.23	0.03 – 6.85	3
Brain and Other CNS	0.0	*	5.8	6.8	6.4	0.00	0.00 – 0.00	1
Breast (Female)	<1	*	122.2	120.4	123.8	0.39	0.01 – 2.15	1
Cervix (Female)	0.0	*	8.0	8.0	8.2	0.00	0.00 – 0.00	1
Colon & Rectum	<1	*	51.1	52.1	49.1	0.53	0.01 – 2.93	1
Corpus Uterus (Female)	<1	*	29.9	26.4	23.3	2.09	0.05 – 11.7	4
Esophagus	0.0	*	5.7	5.5	4.5	0.00	0.00 – 0.00	1
Hodgkin's Lymphoma	0.0	*	2.9	3.0	2.8	0.00	0.00 – 0.00	1
Kidney & Renal Pelvis	<1	*	14.1	14.1	13.6	1.92	0.05 – 10.7	4
Larynx	<1	*	4.7	4.4	3.5	7.61	0.19 – 42.4	4
Leukemia	0.0	*	11.8	11.1	12.2	0.00	0.00 – 0.00	1
Liver & Intrahepatic Bile Duct	0.0	*	5.6	4.3	6.6	0.00	0.00 – 0.00	1
Lung & Bronchus	0.0	*	70.7	75.0	63.1	0.41	0.01 – 2.28	1
Melanoma of the Skin	0.0	*	15.3	17.6	19.6	0.00	0.00 – 0.00	1
Multiple Myeloma	0.0	*	6.0	5.1	5.6	0.00	0.00 – 0.00	1
Non-Hodgkin's Lymphoma	0.0	*	19.1	19.2	19.5	0.00	0.00 – 0.00	1
Oral Cavity & Pharynx	0.0	*	10.4	9.6	10.4	0.00	0.00 – 0.00	1
Ovary	<1	*	12.3	12.3	13.1	3.63	0.09 – 20.2	4
Pancreas	<1	*	12.3	10.9	11.7	4.46	0.54 – 16.1	4
Prostate (Male)	0.0	*	157.5	144.7	159.3	0.37	0.01 – 2.09	1
Stomach	0.0	*	8.1	6.1	7.9	0.00	0.00 – 0.00	1
Testis (Male)	0.0	*	4.9	5.5	5.4	0.00	0.00 – 0.00	1
Thyroid	0.0	*	7.9	8.3	9.6	0.00	0.00 – 0.00	1

Peer Comparison Neighborhoods/Municipalities: Corlett, Downtown, Edgewater, Euclid, Euclid-Green, Highland Hills, Industrial Valley, Linndale, North Collinwood, North Randall, Puritas-Longmead, Riverside, West Boulevard, Woodmere

¹Refer to the methods/limitations section for interpretation of SIRs and Confidence Intervals.

*Rates were suppressed if there were less than 5 cases during 2002-2006. Additionally, to help maintain confidentiality, rates were suppressed if the denominator minus the numerator was less than ten. Rates may also be unstable when case counts are less than 16, thus they should be interpreted with caution in this instance. See the methods/limitations section for additional details.